

Relating Conflict-free Stable Transition and Event Models via Redex Families [★]

Zurab Khasidashvili

*Department of Mathematics and Computer Science
Bar-Ilan University, Ramat-Gan 52900, Israel
khasidz@cs.biu.ac.il*

John Glauert

*School of Information Systems
University of East Anglia, Norwich NR4 7TJ, UK
jrwg@sys.uea.ac.uk*

Abstract

We describe an event-style (or poset) semantics for conflict-free rewrite systems, including term and graph rewriting (possibly with bound variables), the λ -calculus, and other stable transition systems with a residual relation. Our interpretation is based on considering redex-families as events. It treats permutation-equivalent reductions as representing the same concurrent computation. Due to erasure of redexes, Event Structures are inadequate for such an interpretation. We therefore extend the Prime Event Structure model in two different but equivalent ways: by axiomatizing permutation-equivalence on finite configurations, and by axiomatizing the erasure of events, for the conflict-free case, and show that these extended models are equivalent to stable transition models with axiomatized residual and family relations. We then construct finitary prime algebraic domains from the set of configurations in these extended models by defining orderings relative to stable sets of ‘results’. All useful sets of results for which the normalization (by neededness) theorem can be proved are stable.

Keywords: Abstract reduction systems, redex families, event structures, distributive domains.

[★] Part of this work was supported by the Engineering and Physical Sciences Research Council of Great Britain under grant GR/H 41300.

1 Introduction

The primary goal of this paper is to provide a fully adequate event-style concurrent semantics for orthogonal rewrite systems. Event Structures, developed by Winskel, Nielsen and Plotkin [36,32,37,38], are a commonly accepted model of concurrency. Conflict-free Prime Event Structures (PESs) are sufficient for our purposes. A (conflict-free) PES is simply a set of events partially ordered by a *causal dependency* relation (a *poset*) where an event can only dominate a finite number of events (the so called *axiom of finite causes*). The idea is to view a process as performing events (some *atomic* tasks), and an event can occur, only once, after all events on which it depends. Causally independent events are *concurrent* events and can be evaluated in parallel. Stages of a computation are thus represented by *configurations* – left-closed event sets w.r.t. the causal order, and the same configuration represents all sequential computations executing events in that configuration, ignoring the order in which the concurrent events occur.

In the theory of orthogonal rewrite systems, such as the λ -calculus, there is the well established concept of ‘redex-family’, due to Lévy [26,27], which formalizes the idea of the ‘same atomic task’. Therefore it is natural to base our interpretation on considering redex-families as events, and to interpret complete family-reductions (contracting entire families as multi-steps) as configurations of the corresponding PES. This is also justified by the fact that reductions in optimal graph-implementations of orthogonal systems correspond (up to book-keeping steps) to complete family-reductions in the original system [28,23,15,1].¹

Recent advances in the abstract study of the syntax and operational semantics of rewrite and transition systems [35,12,29,11,17,18,20] allow us to address the problem in an entirely abstract setting. Stable Deterministic Residual Structures (SDRSs) and Deterministic Family Structures (DFSs) model computation in orthogonal rewrite systems. DRSs are Abstract Rewrite Systems with an axiomatized residual relation; DFSs are DRSs where in addition the concept of redex-family is axiomatized. Stable DRSs allow [11,17] proofs of analogs of the Normalization and Standardization Theorems [3]. In DFSs one can further prove the Optimality Theorem and the Unique Families Lemma [11]. The latter states that any family can be contracted at most once in a complete family-reduction. It corresponds to the fact that any event may occur at most once in the course of a computation, and is needed for the event structure interpretation of SDRSs based on interpreting families as events.

Thus we can start our interpretation by assuming that an orthogonal rewrite

¹ The book-keeping steps can be expensive in the case of λ -calculus [2].

system is given in the form of a DFS. Furthermore, it has been shown in [20] that any DFS can be given an interpretation by a non-duplicating or affine DFS with zig-zag as the family relation (AZDFS), called the *implementation* of the original DFS: the reduction steps in the implementation DFS correspond to complete family-reduction multi-steps of the original DFS. Therefore it is enough to concentrate on defining PES semantics for AZDFSs.

To view a process as the domain of all its configurations, an *adequacy* requirement is desirable. Namely, that *runs* of a finite configuration (i.e., all its possible sequentializations) should be in one-to-one correspondence with all finite reductions in the equivalence class which the configuration is supposed to represent. Because of the huge importance of *permutation* or *Lévy equivalence* to the whole theory of (not only orthogonal) rewrite systems [27,14,5], and to concurrency theory as well [35], we want to be able to treat permutation-equivalent finite reductions as representing the same concurrent computation, as indeed permutation-equivalent finite reductions result one from another by permuting concurrent consecutive steps. However, this is clearly impossible in the present PES framework [37], as a finite configuration in a PES has only a finite number of runs while, a permutation-equivalence class of a finite reduction may have an infinite number of elements.

For example, consider the λ -term $t = Kx\Omega$, where $K = \lambda xy.x$ and $\Omega = (\lambda x.x)\lambda x.x$. Intuitively, all reductions of the form $t \xrightarrow{\Omega} t \xrightarrow{\Omega} \dots \xrightarrow{\Omega} t \xrightarrow{K} x$, or $\Omega^m K$ in short, consisting of a number of Ω -steps followed by a K -step, are ‘equal’, as the Ω -steps are performed in an erased part of t (or in ‘garbage’). And indeed, permutation equivalence equates all the above reductions. However, there is no means in PESs to equate the corresponding configurations.

The same problem can also be observed from a different angle. In order for an equivalence relation to satisfy the adequacy requirement, the corresponding equivalence classes of reductions must be partially ordered so as to form finitary prime algebraic (or equivalently, distributive) complete lattices (see Definition 55), since the domains of configurations of PESs ordered by inclusion are such lattices [36,37]. However, it is well known that Lévy’s reduction space – permutation equivalence classes of co-initial finite reductions ordered by Lévy’s embedding relation \trianglelefteq (which generates permutation equivalence) – is not in general a lattice but only an upper semi-lattice, as the greatest lower bound of two co-initial finite reductions w.r.t. \trianglelefteq need not exist [27].² As a consequence, although the lattice properties can be restored by taking the ideal completion [4], the resulting domain need not be finitary prime algebraic even when the reduction relation is non-duplicating. For example, Lévy’s configuration domain of the K - Ω term above is not finitary as the K -step $Kx\Omega \xrightarrow{K} x$ is a finitary element but it dominates (w.r.t. \trianglelefteq) an infinite number of elements:

² Recursive Program Schemes are an exception [4].

reductions of the form Ω^m .

Several authors have proposed different ways to get round this problem. Lanve [24] introduced *distributive* equivalence on β -reductions which only allows permutation of steps that cannot erase or duplicate one another, and consequently all equivalent sequences have the same length. Similarly, Corradini et al [8] base their interpretation on the equivalence relation generated by permutations of disjoint redexes only, in a general categorical model of rewriting. Kennaway et al [16] do work with permutation equivalence for orthogonal Term Graph Rewriting, but they only restrict themselves to *needed* events of a normalizable term to cope with problems with erasure. As needed redexes cannot be erased, all needed normalizing reductions are of the same length, and on needed reductions, the permutation and distributive equivalences coincide (since the system is non-duplicating). Clark and Kennaway [7] also work with permutation-equivalence and allow for (finitely) non-normalizable terms, but restrict themselves to *standard* reductions, and their ordering on standard reductions is a strong restriction of Lévy’s embedding relation. And finally, extending work of Boudol [5], Melliès [30] gives a construction of distributive domains from *external* reductions [14], in an axiomatic framework of rewriting. Other authors (e.g., [6,34]) work with linear rewrite/transition systems (no duplication, no erasure), and do not therefore encounter the erasure problem.

In order to cope with the adequacy problem, we extend the PES model in two different but equivalent ways: by axiomatizing permutation-equivalence on event configurations, by directly constraining the class of all equivalence relations on configurations; and by axiomatizing the *inessentiality* or *erasure* relation expressing redundancy of an event for a configuration. The latter extends by further axioms an incomplete axiomatization of the inessentiality relation in [11]. In the resulting extended models, Deterministic *Permutation* and *Erasure* Event Structures, DPESs and DEESs, we can identify finite configurations with different numbers of events but still representing the same permutation-equivalence class of finite reductions. We show that Lévy’s configuration domain – appropriately ordered permutation-equivalence classes of finite configurations in a DEES/DPES – is isomorphic to Lévy’s reduction space of the corresponding AZDFS, and therefore DPESs/DEESs give a *fully adequate* semantics to AZDFSs.

As a consequence, Lévy’s configuration domains in DEESs/DPESs need not be finitary prime algebraic. However, having axiomatized the erasure relation and permutation equivalence on event configurations, we can define orderings on configurations reflecting the growth of information relative to *stable* sets of ‘results’, called *stable orderings*, and thereby reconstruct finitary prime algebraic domains from configuration domains of DEESs/DPESs: The isomorphism between AZDFSs and DEESs/DPESs discussed above induces a

concept of stable sets of configurations in DEESs/DPESs, and the theory of normalization by neededness relative to stable sets of terms in AZDFSs [11] has its counterpart in DEESs/DPESs; and this relativized concept of neededness enables us to show that the stable orderings on configuration domains of DEESs/DPESs do form finitary prime algebraic domains.

The paper is organized as follows. In the next section, we review the existing theory of DRSs and DFSs used in this paper. In Section 3 we introduce DPESs and DEESs and establish their equivalence. In Section 4 we relate the latter event models with affine SDRSs (ASDRSs) and AZDFSs, showing the equivalence of the four models. In Section 5, we define Lévy’s configuration domain in a DPES/DEES and show its isomorphism with Lévy’s reduction space of the corresponding ASDRS. We study normalization by neededness in DEESs in Section 6, and use it to construct finitary prime algebraic domains from configurations in DEESs/DPESs in Section 7. Conclusions appear in Section 8.

2 Deterministic Residual and Family Structures

In this section, we recall some basic theory of *Deterministic Residual* and *Family Structures* (DRSs and DFSs), developed in [11,17,18,20]. DRSs and DFSs are *Abstract Reduction Systems* (ARSs) with an axiomatized notion of residual. In DFSs, in addition, Lévy’s concept of redex-family is axiomatized. Related abstract residual models are studied in [13,35,12,29].

Our definition of ARSs allows multiple transitions with the same source and target, unlike Klop’s definition [22].

Definition 1 An ARS is a triple $A = (Ter, Red, \rightarrow)$ where Ter is a set of *terms*, ranged over by t, s, o, e ; Red is a set of *redexes* (or *redex occurrences*), ranged over by u, v, w ; and $\rightarrow: Red \mapsto (Ter \times Ter)$ is a function such that for any $t \in Ter$ there is only a finite set of $u \in Red$ such that $\mapsto(u) = (t, s)$, written $t \xrightarrow{u} s$. This set will be known as the redexes of term t , where $u \subseteq t$ denotes that u is a member of the redexes of t and $U \subseteq t$ denotes that U is a subset of the redexes. Note that \rightarrow is a *total* function, so one can identify u with the triple $t \xrightarrow{u} s$. A *reduction* is a sequence $t \xrightarrow{u_1} t_2 \xrightarrow{u_2} \dots$. Reductions are denoted by P, Q, N . We write $P : t \twoheadrightarrow s$ or $t \xrightarrow{P} s$ if P denotes a reduction (sequence) from t to s . If P is finite and its final term coincides with the initial term of Q , then $P + Q$ denotes the concatenation of P and Q . Finally, u also denotes the reduction that contracts u .

Definition 2 (Deterministic Residual Structure [11]) A *Deterministic Residual Structure* (DRS) is a pair $\mathcal{R} = (A, /)$, where A is an ARS and $/$ is

a *residual* relation on redexes relating redexes in the source and target term of every reduction $t \xrightarrow{u} s \in A$, such that for $v \subseteq t$, the set v/u of *residuals of v under u* is a set of redexes of s ; a redex in s may be a residual of only one redex in t under u , and $u/u = \emptyset$. If v has more than one u -residual, then u *duplicates* v . If $v/u = \emptyset$, then u *erases* v . A redex of s which is not a residual of any $v \subseteq t$ under u is said to be *u -new* or *created* by u . The set u/P of residuals of u under any finite reduction P is defined by transitivity.

A *development* of $U \subseteq t$ is a reduction $P : t \rightarrow$ that only contracts residuals of redexes from U ; it is *complete* if it is finite and $U/P = \cup_{u \in U} u/P = \emptyset$. Development of \emptyset is identified with the empty reduction. U will also denote a complete development of $U \subseteq t$. The residual relation satisfies the following two axioms:

[FD] (*Finite Developments* [12]) All developments are terminating; all complete developments of $U \subseteq t$ end at the same term; and residuals of a redex $v \subseteq t$ under all complete developments of U are the same.

[weak acyclicity] ([35]) Let $u, v \subseteq t$, $u \neq v$, and $u/v = \emptyset$. Then $v/u \neq \emptyset$.³

We call a DRS \mathcal{R} *stable* (SDRS) if the following axiom is satisfied:

[stability] If $u, v \subseteq t$ are different redexes, $t \xrightarrow{u} e$, $t \xrightarrow{v} s$, and u creates a redex $w \subseteq e$, then the redexes in $w/(v/u)$ are not u/v -residuals of redexes of s , i.e., they are created along u/v .

$$\begin{array}{ccccc}
 t & \xrightarrow{u} & e & \xrightarrow{w} & \\
 v \downarrow & & \downarrow v/u & & \\
 s & \xrightarrow{u/v} & o & \xrightarrow{w/(v/u)} &
 \end{array}$$

We call an SDRS *non-duplicating* or *affine*, ASDRS, if its residual relation is non-duplicating. Note that, since the only observables of DRSs are redexes, duplicating syntactic rewrite systems may still form ASDRSs. For example, the DRS corresponding to innermost reductions in an orthogonal TRS is an ASDRS, although innermost redexes may duplicate their arguments.

In a DRS \mathcal{R} , the residual relation on redexes extends to all co-initial reductions exactly as in syntactic orthogonal rewrite systems [14,26,27,35]: $(P_1 + P_2)/Q = P_1/Q + P_2/(Q/P_1)$ and $P/(Q_1 + Q_2) = (P/Q_1)/Q_2$, and *Lévy-equivalence* or *permutation-equivalence* is defined as the smallest relation on co-initial reductions satisfying: $U + V/U \approx V + U/V$ and $Q \approx Q' \Rightarrow P + Q + N \approx P + Q' + N$, where U and V are complete developments of redex sets in the same term. Further, one defines $P \trianglelefteq Q$ iff $P/Q = \emptyset$, and can show that $P \approx Q$ iff $P \trianglelefteq Q$ and $Q \trianglelefteq P$; and $P \trianglelefteq Q$ iff $Q \approx P + N$ for some N . Intuitively, $P \trianglelefteq Q$ expresses that


³ This axiom is called [acyclicity] in [11], and is axiom (4) in [35].

Q does more work than P , and Q/P is the part of Q that remains from it after P . Finally, one shows that for any co-initial reductions P, Q , $P \sqcup Q \approx Q \sqcup P$, where $P \sqcup Q = P + Q/P$, and is the greatest lower bound of P and Q in the reduction space ordered by Lévy's embedding relation \trianglelefteq . The above relations can equivalently be defined also using Klop's method of commutative diagrams [21,3]. Finally, we will need the following *Cube Lemma*: for any finite co-initial reductions P, Q and N , $N/(P \sqcup Q) = N/(Q \sqcup P)$.

The following two lemmas, established in [11,17], are fundamental in proving properties in SDRSs. They extend [weak acyclicity] and [stability] axioms from one step reductions to any co-initial *external* reductions, that is, reductions that do not contract redexes having common residuals.

Definition 3 • Let $u \in U \subseteq t$ and $P : t \rightarrow o$. We call P *external* to U (resp. u) if P does not contract residuals of redexes in U (resp. residuals of u).

- Let $P : t_0 \xrightarrow{P_i} t_i \xrightarrow{u_i} t_{i+1} \rightarrow t_n$ and $Q : t_0 = s_0 \xrightarrow{Q_j} s_j \xrightarrow{v_j} s_{j+1} \rightarrow s_m$. Let $U_{i,j} = u_i/(Q_j/P_i)$ and $V_{i,j} = v_j/(P_i/Q_j)$ (see diagram). We call P *external* to Q if for any i, j , $U_{i,j} \cap V_{i,j} = \emptyset$.



Lemma 4 (Stability) Let $P : t \rightarrow s$ be external to $Q : t \rightarrow e$, in a stable DRS, and let P create redexes $W \subseteq s$. Then the residuals $W/(Q/P)$ of redexes in W are created by P/Q , and Q/P is external to W .

Lemma 5 (Weak Acyclicity) Let P, Q be co-initial finite reductions in a DRS, and let P be external to Q . Then $Q \not\approx P$.

Next we recall some definitions and results from [17] concerning standardization of reductions in ASDRSs, used in this paper.

Definition 6 • Let $P : t \rightarrow o$ and $u \subseteq t$, in a DRS. We call u *erased* in P or *P -erased* if $u/P = \emptyset$. We say that P *discards* u if P is external to u and erases it.

- We call u *P -needed* if there is no $Q \approx P$ that is external to u , and call it *P -unnneeded* otherwise. We call u *P -essential* if there is no $Q \approx P$ that discards u , and *P -inessential* otherwise.

We extend these concepts to reductions co-initial with those containing

u as a redex of one of its terms.

- Let $Q : t \rightarrow o$, $P : t \xrightarrow{P'} s \rightarrow e$, and $u \subseteq s$. We say that u is Q -needed, or more precisely, $P'u$ is Q -needed, if u is Q/P' -needed. We call P Q -needed if every redex contracted in P is. We call P *self-needed* if it is P -needed. The other concepts above are extended in the same way.

Note that P -neededness, P -erasure, and P -essentiality do not depend on the choice of a reduction in the class $\langle P \rangle_L$ of reductions Lévy-equivalent to P , since $u/P = u/Q$ when $P \approx Q$, by the Cube Lemma. The *discards* concept however does depend on the particular reduction in the permutation-equivalence class.

Lemma 7 Let $P : s \rightarrow t \xrightarrow{u} e \rightarrow o$ in an ASDRS.

- (1) $w \in t$ is P -needed iff it is P -erased and P -essential.
- (2) If $P : t \rightarrow s' \xrightarrow{w} o$, then $w \in s'$ is P -needed.
- (3) If u creates $v \subseteq e$ and u is P -unneeded (resp. P -inessential), then so is v .
- (4) If $u \neq v \subseteq t$, then v is P -needed (P -essential) iff v has a P -needed (P -essential) residual in e .

Self-needed reductions play the role of *standard* reductions in SDRSs, and the following algorithm is a standardization procedure for reductions in ASDRSs. Let $P : t \rightarrow s$. The *canonical standard variant of P* , $ST(P)$, is defined as follows: If $P = \emptyset$, then $ST(P) = \emptyset$. Otherwise, let $v \subseteq t$ be such that it is P -needed and its residual is contracted in P first among P -needed residuals of P -needed redexes in t (existence of such v follows from Lemma 7). Then $ST(P) = v + ST(P/v)$. We write $Q \in STV(P)$ if $Q \in STA$ and $Q \approx P$, where STA denotes the set of all standard reduction, and call Q a *standard variant* of P . In particular, $ST(P) \in STV(P)$.

Theorem 8 (Standardization) For any finite reduction P in a stable non-duplicating DRS, $ST(P)$ is a standard reduction Lévy-equivalent to P .

It is shown in [18,20] that all standard variants of a finite reduction P can be constructed effectively.

Definition 9 (Deterministic Family Structure [11]) A *Deterministic Family Structure* (DFS) is a triple $\mathcal{F} = (\mathcal{R}, \simeq, \hookrightarrow)$, where \mathcal{R} is a DRS; \simeq is an equivalence relation on redexes with *histories*; and \hookrightarrow is the *contribution* relation on co-initial families, defined as follows:

- (1) For any co-initial finite reductions P and Q , a redex Qv in the final term of Q (read as v with history Q) is called a *copy* of a redex Pu if $P \triangleleft Q$, i.e., $P + Q/P \approx Q$, and v is a Q/P -residual of u ; the *zig-zag* relation \simeq_z is the symmetric and transitive closure of the copy relation. The *family* relation \simeq is an equivalence relation among redexes with histories containing \simeq_z . A *family* is an equivalence class of the family relation; families are ranged

over by ϕ, ψ, \dots . $Fam(\)$ denotes the family of its argument.

(2) Further, \simeq and \hookrightarrow satisfy the following axioms:

[initial] Let $u, v \subseteq t$ and $u \neq v$, in \mathcal{R} . Then $Fam(\emptyset_t u) \neq Fam(\emptyset_t v)$, where \emptyset_t is the empty reduction starting from t .

[contribution] $\phi \hookrightarrow \phi'$ iff for any $Pu \in \phi'$, P contracts at least one redex in ϕ .

[creation] Let $e \xrightarrow{P} t \xrightarrow{u} s$, and let u create $v \subseteq s$. Then $Fam(Pu) \hookrightarrow Fam((P + u)v)$.

[FFD] (*Finite Family Developments*) Any reduction that contracts redexes of a finite number of families is terminating.⁴

Note that [contribution] can be viewed as a definition of \hookrightarrow rather than as an axiom. It is shown in [11] that every DFS is a stable DRS.

Convention In this paper we only consider *comma-DRSs* and *comma-DFSs*. That is, the term set of any DRS will be the *reduction graph* of a term, called the *initial* term, i.e., the set of terms to which the initial term is reducible. The initial term will often be denoted by t_\emptyset , by analogy with the initial state \emptyset in an Event Structure. Further, in DFSs, families will always be considered relative to t_\emptyset , i.e., all histories start with t_\emptyset . Unless otherwise stated, by a *reduction* we mean a *finite* reduction. Reductions starting from t_\emptyset will also be called *initial* reductions.

Definition 10 • We call a DFS \mathcal{F} a *zig-zag* DFS, ZDFS, if its family relation is the zig-zag \simeq_z . Affine ZDFSs will be called AZDFSs.

- We call an affine DFS *separable* if, for any redex Pv , v cannot create two different redexes in the same family. That is, if v creates w', w'' and $w' \neq w''$, then $Fam((P + v)w') \neq Fam((P + v)w'')$.

The following lemma from [18,20] gives a useful characterization of \trianglelefteq and \approx in ASDRSs via zig-zag classes. Below, $FAM(P)$ (resp. $SFAM(P)$) denotes the set of *zig-zag classes* whose member (resp. P -needed) redexes are contracted in P , in an ASDRS.

Proposition 11 Let P and Q be initial reductions in an AZDFS. Then $P \trianglelefteq Q$ iff $\exists Q' \approx Q : FAM(P) \subseteq FAM(Q')$, and $P \approx Q$ iff $SFAM(P) = SFAM(Q)$. Further, $SFAM(P) = FAM(ST(P)) = \bigcap_{N \approx P} FAM(N)$. Consequently, if $P, Q \in STA$, then $P \approx Q$ iff $FAM(P) = FAM(Q)$ (since in that case $P = ST(P)$ and $Q = ST(Q)$).

The results of this paper are based on the following properties of ASDRSs, which show that ASDRSs and AZDFSs are essentially the same transition models, and that every zig-zag family has a unique (up to an equivalence on histories) minimal element [18,20].

⁴ This axiom is called [termination] in [11].

Theorem 12 (Affine Families) Let \mathcal{R} be an ASDRS. Then $\mathcal{F}_R = (\mathcal{R}, \simeq_z, \hookrightarrow_z)$ is an affine zig-zag DFS, called the *corresponding* ZDFS of \mathcal{R} . Further, \mathcal{F}_R is the only separable DFS with R as the underlying SDRS, and in general, an affine DFS is separable iff it is a zig-zag DFS.

Theorem 13 (Affine Extraction) Every redex Pv in an ASDRS \mathcal{R} has a zig-zag related redex $P'v'$ in *canonical* form, meaning that $P' \in STA$, its last step creates v' , and for any $P''v'' \simeq_z Pv$, $\exists N' : P'' \approx P' + N' \wedge v'' = v'/N'$. For any other canonical form P^*v^* of Pv , $P' \approx P^*$ and $v' = v^*$. Further, P' contracts exactly one redex in every contributor zig-zag family of Pv , in the corresponding AZDFS $\mathcal{F}_R = (\mathcal{R}, \simeq_z, \hookrightarrow_z)$ of \mathcal{R} .

3 Deterministic Permutation and Erasure Event Structures

In this section, we introduce Deterministic Permutation and Erasure Event Structures, DPESs and DEESs, which extend conflict-free Prime Event Structures by an axiomatized permutation-equivalence and erasure relations, respectively, and establish an equivalence between them.

The purpose of this extension is to capture *directly* the phenomenon of *erasure* in event/poset models. The phenomenon is typical for languages based on rewrite systems, but is not confined to them. In higher-order process calculi (e.g., [33]) processes (that can fire events) can be passed around as messages and therefore erased. Event Structures are linear in nature and consequently the configuration domains enjoy very nice lattice properties; therefore Event Structures have been very successful in modelling *linear* process calculi where no actions can be erased or duplicated (see e.g., [6,37]), such as CCS [31]. However, any attempt to adequately interpret higher-order process languages in Event Structures would face the same erasure problems discussed in the introduction. As in the case of reductions, the aim of axiomatizing permutation equivalence on event configurations is to be able to equate configurations which represent the same concurrent computation and differ only by irrelevant or *inessential* events. We believe that any equivalence relation on configurations fully capturing concurrency should contain the permutation-equivalence.

A *Prime Event Structure* (PES) [36] is a triple $\mathcal{E} = (E, Con, \leq)$, where E is a set of *events*, ranged over by e, e_1, \dots ; the *consistency predicate* Con is a non-empty set of finite subsets of E , denoted by X, Y, \dots ; and the *causal dependency relation* \leq is a partial order on E , such that $\{e\} \in Con$, $Y \subseteq X \in Con \Rightarrow Y \in Con$, $X \in Con \wedge \exists e' \in X. e \leq e' \Rightarrow X \cup \{e\} \in Con$, and $\{e' \mid e' \leq e\}$ is finite for any $e \in E$.

We only consider *conflict-free* or *deterministic* PESs, where no event can pre-

vent others from occurring, and therefore the consistency predicate is the set of all finite subsets of E and will be omitted. Finite *configurations* of \mathcal{E} are finite *left-closed subsets* α, β, \dots of E , i.e., subsets $\mathcal{L}_{fin}(E) = \{\alpha \subseteq_{fin} E \mid e \in \alpha \wedge e' < e \Rightarrow e' \in \alpha\}$. Except for Section 7, we only consider finite configurations, and will omit ‘finite’.

Below, for better readability, we write $\alpha + e$ for $\alpha \cup \{e\}$, and write $\alpha - e$ for $\alpha \setminus \{e\}$. Further, we define $\lceil e \rceil^{\leq} = \{e' \mid e' \leq e\}$, $\lceil e \rceil^{<} = \{e' \mid e' < e\}$, $\lceil e \rceil^{\geq} = \{e' \mid e \geq e'\}$ and $\lceil e \rceil^{>} = \{e' \mid e' > e\}$.

Definition 14 A *Deterministic Permutation Event Structure* (DPES) is a triple $\mathcal{P} = (E, \leq, \approx)$, where $\mathcal{E} = (E, \leq)$ is a conflict-free PES and $\approx \subseteq \mathcal{L}_{fin}(E) \times \mathcal{L}_{fin}(E)$ is an equivalence relation, called *Lévy- or permutation-equivalence*, satisfying the following axioms, where $\alpha, \beta, \gamma \in \mathcal{L}_{fin}(E)$:

- [P0] $\forall e \in E : \lceil e \rceil^{<} \not\approx \lceil e \rceil^{\leq}$;
- [P1] $\alpha \approx \beta \wedge \alpha \cap \beta \subseteq \gamma \subseteq \alpha \cup \beta \Rightarrow \alpha \approx \gamma$;
- [P2] $\alpha \approx \beta \wedge \alpha + e, \beta + e \in \mathcal{L}_{fin}(E) \Rightarrow \alpha + e \approx \beta + e$;
- [P3] $\alpha \approx \alpha - e \wedge e < e' \wedge \alpha + e' \in \mathcal{L}_{fin}(E) \Rightarrow \alpha \approx \alpha + e'$;

DEESs extend *Deterministic Computation Structures* [11] by further erasure axioms to enable a fully adequate treatment of erasure.

Definition 15 A *Deterministic Erasure Event Structure* (DEES) is a triple $\mathcal{C} = (E, \leq, \triangleright)$, where $\mathcal{E} = (E, \leq)$ is a conflict-free PES and $\triangleright \subseteq \mathcal{L}_{fin}(E) \times E$ is *inessentiality* or *erasure* relation (read $\alpha \triangleright e$ as: ‘ e is α -inessential’), satisfying the following axioms, where $\alpha, \beta \in \mathcal{L}_{fin}(E)$:

- [E0] $\forall e \in E : \emptyset \not\triangleright e$;
- [E1] $\alpha \triangleright e \wedge \alpha \subseteq \beta \in \mathcal{L}_{fin}(E) \Rightarrow \beta \triangleright e$;
- [E2] $\alpha \triangleright e' \wedge \alpha \triangleright e \wedge \alpha - e' \in \mathcal{L}_{fin}(E) \Rightarrow \alpha - e' \triangleright e$;
- [E3] $\alpha \triangleright e \wedge e < e' \Rightarrow \alpha \triangleright e'$;
- [E4] $\alpha \cup \lceil e \rceil^{<} \triangleright e \Rightarrow \alpha \triangleright e$.

We may write $\alpha \triangleright X$ if $\forall e \in X : \alpha \triangleright e$. Sometimes we write $\mathcal{L}_{fin}(\mathcal{P})$ for $\mathcal{L}_{fin}(E)$ if E is the domain of a DPES \mathcal{P} , and similarly for $\mathcal{L}_{fin}(\mathcal{C})$.

Before explaining the intuition behind the axioms, we first define translations between DPESs and DEESs.

Definition 16 For a DPES $\mathcal{P} = (E, \leq, \approx)$, we define an *erasure* relation $\triangleright_{\approx} \subseteq \mathcal{L}_{fin}(E) \times E$ and the corresponding structure $\mathcal{C}_{\mathcal{P}} = (E, \leq, \triangleright_{\approx})$ as follows:

- $\alpha \triangleright_{\approx} e$ iff $\alpha \cup \lceil e \rceil^{\leq} \approx (\alpha \cup \lceil e \rceil^{\leq}) \setminus \lceil e \rceil^{\geq}$.

In particular, when $e \in \alpha$ and $\alpha - e \in \mathcal{L}_{fin}(E)$,

- $\alpha \triangleright_{\approx} e$ iff $\alpha \approx \alpha - e$.

Definition 17 For a DEES $\mathcal{C} = (E, \leq, \triangleright)$, we define an equivalence relation $\approx_{\triangleright} \subseteq \mathcal{L}_{fin}(E) \times \mathcal{L}_{fin}(E)$ and the corresponding structure $\mathcal{P}_{\mathcal{C}} = (E, \leq, \approx_{\triangleright})$ as follows:

- $\alpha \approx_{\triangleright} \beta$ iff $ST_{\triangleright}(\alpha) = ST_{\triangleright}(\beta)$, where $ST_{\triangleright}(\alpha) = \{e \in \alpha \mid \alpha \not\triangleright e\}$. (The subscript \triangleright in $ST_{\triangleright}(\alpha)$ will often be omitted.) $ST(\alpha)$ will be called the *standard variant* of α .

By looking at a configuration α as an initial reduction P_{α} firing redexes corresponding to events in α , and by interpreting $\alpha \triangleright e$ as ‘ Qv is P -inessential’, where Qv is a redex representing v (so Q contracts redexes corresponding to events on which e causally depends), it is not difficult to understand the intuition behind the erasure axioms. For example, [E0] says that an empty reduction cannot erase any redex, and [E1] says that if Qv is P_{α} -inessential and $P_{\alpha} \leq P_{\beta}$, then Qv is P_{β} -inessential as well. This intuition will become more precise in the next section. Further, a close inspection of the above definitions shows that the DPES axiom [P0] is a counterpart of the DEES axioms [E0] and [E4], and similarly for [P3] and [E3]. The combination of [P1] and [P2] have the same effect as that of [E1] and [E2]. [E4] does not have a ‘direct’ counterpart among DPES axioms, but it corresponds to the fact that, according to Definition 6, a redex with history Pv is Q -inessential iff v is Q/P -inessential. Similarly, [P0] corresponds to Lemma 7.(2).

In order to establish equivalence of DEESs and DPESs, we need a few lemmas.

Lemma 18 Let $\alpha \in \mathcal{L}_{fin}(\mathcal{C})$, in a DEES \mathcal{C} . Then $ST(\alpha) \in \mathcal{L}_{fin}(\mathcal{C})$. Further, $ST(ST(\alpha)) = ST(\alpha)$, hence $\alpha \approx_{\triangleright} ST(\alpha)$.

Proof Let $e \in ST(\alpha) \wedge e' < e$. Then $e \in \alpha$ and $\alpha \not\triangleright e$, implying by [E3] that $\alpha \not\triangleright e'$. Since $\alpha \in \mathcal{L}_{fin}(\mathcal{C})$ and $e' < e$, $e' \in \alpha$, implying $e' \in ST(\alpha)$. Hence $ST(\alpha) \in \mathcal{L}_{fin}(\mathcal{C})$. If, on the contrary, $ST(ST(\alpha)) \neq ST(\alpha)$, i.e., $\exists e'' \in ST(\alpha) : ST(\alpha) \triangleright e''$, then $\alpha \triangleright e''$ by [E1], contradicting $e'' \in ST(\alpha)$. Now $\alpha \approx_{\triangleright} ST(\alpha)$ follows immediately from Definition 17.

Lemma 19 Let $\alpha \in \mathcal{L}_{fin}(\mathcal{C})$, in a DEES \mathcal{C} . Then $\alpha \triangleright e$ iff $ST(\alpha) \triangleright e$.

Proof

(\Rightarrow) Let $\alpha = ST(\alpha) \cup \{e_1, \dots, e_n\}$, where the enumeration e_1, \dots, e_n respects the order on E , let $\alpha_0 = \alpha$, and let $\alpha_{i+1} = \alpha_i + e_{i+1}$ (hence $\alpha = \alpha_n$). Denote $e = e_0$. By the definition of $ST(\alpha)$ and the assumption, we have $\alpha_n \triangleright e_i$ for all $i = 0, 1, \dots, n$. By [E2], we have from $\alpha_n \triangleright e_n$ and $\alpha_n \triangleright e_i$ that $\alpha_{n-1} \triangleright e_i$. Hence, again by [E2], $\alpha_{n-2} \triangleright e_i$, and so on. Finally, we get $\alpha_0 \triangleright e_i$ for all i , and in particular, $\alpha_0 \triangleright e$.

(\Leftarrow) By [E1].

Lemma 20 Let $\alpha, \beta \in \mathcal{L}_{fin}(E)$ and $\alpha \subseteq \beta$, in a DEES \mathcal{C} . Then $ST(\alpha) = ST(\beta)$ iff $\beta \triangleright \beta \setminus \alpha$.

Proof

(\Rightarrow) By Definition 17, since $\beta \setminus \alpha \subseteq \beta \setminus ST(\alpha)$.

(\Leftarrow) Since $\beta \triangleright \beta \setminus \alpha$, $ST(\beta) \subseteq \alpha$. Let $e' \in ST(\beta)$. Then $e' \in \beta$ and $\beta \not\triangleright e'$, implying by [E1] and $\alpha \subseteq \beta$ that $\alpha \not\triangleright e'$, i.e., $e' \in ST(\alpha)$. Hence $ST(\beta) \subseteq ST(\alpha)$. For the converse, take $e'' \in ST(\alpha)$. Then $e'' \in \alpha$ and $\alpha \not\triangleright e''$. Therefore, $e'' \in \beta$ and by [E1] $ST(\beta) \not\triangleright e''$, implying by Lemma 19 that $\beta \not\triangleright e''$, i.e., $e'' \in ST(\beta)$. Hence $ST(\alpha) = ST(\beta)$.

Proposition 21 Let $\alpha, \beta, \gamma \in \mathcal{L}_{fin}(\mathcal{C})$, in a DEES \mathcal{C} , and let $\alpha \cap \beta \subseteq \gamma \subseteq \alpha \cup \beta$. Then $ST(\alpha) = ST(\beta) \Leftrightarrow \gamma \triangleright \alpha \cup \beta \setminus \alpha \cap \beta$. Consequently, $\alpha \approx_{\triangleright} \beta$ iff $\gamma \triangleright \alpha \cup \beta \setminus \alpha \cap \beta$.

Proof

(\Rightarrow) $ST(\alpha) = ST(\beta) \Rightarrow ST(\alpha) \subseteq \alpha \cap \beta$, and by the definition of $ST(\alpha)$, Lemma 19, and [E1], $ST(\alpha) = ST(\beta) \Rightarrow ST(\alpha) \triangleright \beta \setminus \alpha \cup \alpha \setminus \beta \Rightarrow \gamma \triangleright \alpha \cup \beta \setminus \alpha \cap \beta$.

(\Leftarrow) By [E1] and Lemma 20, $\alpha \cap \beta \triangleright \alpha \cup \beta \setminus \alpha \cap \beta \Rightarrow \alpha \triangleright \alpha \setminus \beta \wedge \beta \triangleright \beta \setminus \alpha \Rightarrow ST(\alpha) = ST(\alpha \cap \beta) = ST(\beta)$.

Theorem 22 For any DEES $\mathcal{C} = (E, \leq, \triangleright)$, $\mathcal{P}_{\mathcal{C}} = (E, \leq, \approx_{\triangleright})$ is a DPES.

Proof We need to show that \approx_{\triangleright} satisfies the permutation axioms [P0] – [P3].

[P0]: Let $e \in E$. Note that $[e]^{\leq} \triangleright e$ would imply by [E2] that $[e]^{\leq} \triangleleft e$, implying $\emptyset \triangleright e$ by [E4], contradicting [E0]. Hence $[e]^{\leq} \not\triangleright e$, i.e., $e \in ST([e]^{\leq})$, and therefore $ST([e]^{\leq}) \neq ST([e]^{\triangleleft})$. Thus $[e]^{\leq} \not\approx_{\triangleright} [e]^{\triangleleft}$ by Definition 17.

[P1]: Let $\alpha \approx_{\triangleright} \beta$, $\alpha \cap \beta \subseteq \gamma \subseteq \alpha \cup \beta$, and $\gamma \in \mathcal{L}_{fin}(E)$. By Definition 17, $ST(\alpha) = ST(\beta)$, hence $ST(\alpha) \subseteq \alpha \cap \beta \subseteq \gamma$. Further, $\forall e \in \alpha \cup \beta \setminus ST(\alpha)$, $\alpha \triangleright e$ or $\beta \triangleright e$, hence $ST(\alpha) \triangleright e$ by Lemma 19. Hence $ST(ST(\alpha)) = ST(\gamma)$ by Proposition 21, and $ST(\alpha) = ST(\gamma)$ by Lemma 18, implying $\gamma \approx_{\triangleright} \alpha$ by Definition 17.

[P2]: Let $\alpha \approx_{\triangleright} \beta$ and $\alpha + e, \beta + e \in \mathcal{L}_{fin}(E)$. By Definition 17, $ST(\alpha) = ST(\beta)$. Assume first that $ST(\alpha) \triangleright e$. Then $\alpha + e \triangleright e$ by [E1] and $ST(\alpha + e) = ST(\alpha)$ by Proposition 21. Similarly, $ST(\beta + e) = ST(\beta)$. Hence, $ST(\alpha + e) = ST(\beta + e)$, implying by Definition 17 that $\alpha + e \approx_{\triangleright} \beta + e$. Now let $ST(\alpha) \not\triangleright e$. Then, by Lemma 19 and [E3], $\forall e' \in \alpha \setminus ST(\alpha) : e' \not\triangleright e$. Thus $\alpha + e \in \mathcal{L}_{fin}(E) \Rightarrow ST(\alpha) + e \in \mathcal{L}_{fin}(E)$. By Lemma 19, $ST(\alpha) \not\triangleright e$ implies $\alpha \not\triangleright e$, thus $e \in ST(\alpha)$ iff $e \in \alpha$, and therefore $\alpha + e \setminus (ST(\alpha) + e) = \alpha \setminus ST(\alpha)$. Hence $\alpha + e \triangleright \alpha + e \setminus (ST(\alpha) + e)$ by [E1] (as $\alpha \triangleright \alpha \setminus ST(\alpha)$), and we have by

Proposition 21 that $ST(\alpha + e) = ST(ST(\alpha) + e)$, thus $\alpha + e \approx_{\triangleright} ST(\alpha) + e$ by Definition 17. Similarly, $\beta + e \approx_{\triangleright} ST(\beta) + e$, implying $\alpha + e \approx_{\triangleright} \beta + e$ (since $ST(\alpha) = ST(\beta)$).

[P3]: Let $\alpha \approx_{\triangleright} \alpha - e$, $e < e'$ and $\alpha + e' \in \mathcal{L}_{fin}(E)$ (hence $e \in \alpha$). By Definition 17, $ST(\alpha) = ST(\alpha - e)$, implying $\alpha \triangleright e$. Hence, by [E3], $\alpha \triangleright e'$, implying by [E1] that $\alpha + e' \triangleright e'$. Hence, by Proposition 21, $ST(\alpha + e') = ST(\alpha)$, thus $\alpha + e' \approx_{\triangleright} \alpha$ by Definition 17.

Theorem 23 For any DPES $\mathcal{P} = (E, \leq, \approx)$, $\mathcal{C}_P = (E, \leq, \triangleright_{\approx})$ is a DEES.

Proof We need to show that \triangleright_{\approx} satisfies the erasure axioms [E0] – [E4].

[E0]: By Definition 16 and [P0].

[E3]: Let $\alpha \triangleright_{\approx} e$ and $e < e'$. We want to show that $\alpha \triangleright_{\approx} e'$. We can assume that $\nexists e'' : e < e'' < e'$. Assume first that $e, e' \in \alpha$ (hence $\alpha \cup [e]^{\leq} = \alpha \cup [e']^{\leq} = \alpha$). By Definition 16, $\alpha \triangleright_{\approx} e \Rightarrow \alpha \approx \alpha \setminus [e]^{\geq}$. But $\alpha \setminus [e]^{\geq} \subseteq \alpha \setminus [e']^{\geq} \subseteq \alpha$, implying by [P1] that $\alpha \approx \alpha \setminus [e']^{\geq}$, i.e., by Definition 16, $\alpha \triangleright_{\approx} e'$. Now let $e \in \alpha$ and $e' \notin \alpha$. Let $\gamma = \alpha \setminus [e]^{\geq}$. By Definition 16, $\alpha \triangleright_{\approx} e$ iff $\gamma \triangleright_{\approx} e$ iff $\gamma \approx \gamma - e$. Further, $\gamma \cup [e']^{\leq} = \gamma + e'$ and $\gamma \cup [e']^{\leq} \setminus [e']^{\geq} = \gamma$. By [P3], $\gamma \approx \gamma + e'$, implying $\gamma \triangleright_{\approx} e'$ and equivalently $\alpha \triangleright_{\approx} e'$ by Definition 16. Finally, let $e, e' \notin \alpha$. Take $\beta = \alpha \cup [e]^{\leq}$. Then, by Definition 16, $\alpha \triangleright_{\approx} e$ iff $\beta \triangleright_{\approx} e$ and $\alpha \triangleright_{\approx} e'$ iff $\beta \triangleright_{\approx} e'$, and we have $e \in \beta$ and $e' \notin \beta$ like in the previous case, and [E3] follows.

[E1]: Let $\alpha \triangleright_{\approx} e \wedge \alpha \subseteq \beta \in \mathcal{L}_{fin}(E)$. We can assume that $\beta = \alpha + e'$ and $e' \notin \alpha$. If $e \geq e'$, then $\alpha + e' \cup [e]^{\leq} = \alpha \cup [e]^{\leq}$ and $\alpha + e' \cup [e]^{\leq} \setminus [e]^{\geq} = \alpha \cup [e]^{\leq} \setminus [e]^{\geq}$, thus $\alpha \triangleright_{\approx} e$ iff $\alpha + e' \triangleright_{\approx} e$ by Definition 16. If $e \not\geq e'$ and $e' \not\leq e$, then $\alpha + e' \cup [e]^{\leq} = \alpha \cup [e]^{\leq} + e'$ and $\alpha + e' \cup [e]^{\leq} \setminus [e]^{\geq} = \alpha \cup [e]^{\leq} \setminus [e]^{\geq} + e'$, and $\alpha + e' \triangleright_{\approx} e$ follows from $\alpha \triangleright_{\approx} e$ by [P2] and Definition 16. Finally, if $e < e'$, then $\alpha + e' \in \mathcal{L}_{fin}(E) \Rightarrow e \in \alpha$. By [E3] which we have already proven above for \triangleright_{\approx} , $\alpha \triangleright_{\approx} e \Rightarrow \alpha \triangleright_{\approx} e'$; thus by Definition 16 we have from $\alpha \triangleright_{\approx} e$ that $\alpha = \alpha \cup [e]^{\leq} \approx \alpha \cup [e]^{\leq} \setminus [e]^{\geq}$, and we have from from $\alpha \triangleright_{\approx} e'$ that $\alpha \approx \alpha + e'$. Hence $\alpha + e' \cup [e]^{\leq} \setminus [e]^{\geq} \approx$ (since $e < e'$) $\approx \alpha \cup [e]^{\leq} \setminus [e]^{\geq} \approx \alpha \approx \alpha + e' = \alpha + e' \cup [e]^{\leq}$, which means by Definition 16 that $\alpha + e' \triangleright e$.

[E2]: We can assume that $e' \in \alpha$. Let $\alpha \triangleright_{\approx} e'$, $\alpha \triangleright_{\approx} e$ and $\alpha - e' \in \mathcal{L}_{fin}(E)$. By Definition 16, $\alpha \triangleright_{\approx} e \Rightarrow \alpha \cup [e]^{\leq} \approx \alpha \cup [e]^{\leq} \setminus [e]^{\geq}$ and $\alpha \triangleright_{\approx} e' \Rightarrow \alpha \approx \alpha - e'$, implying by repeated application of [P2] that $\alpha \cup [e]^{\leq} \approx \alpha - e' \cup [e]^{\leq}$. It is easy to check that $(\alpha \cup [e]^{\leq} \setminus [e]^{\geq}) \cap (\alpha - e' \cup [e]^{\leq}) = (\alpha \cup [e]^{\leq}) \cap (\alpha - e' \cup [e]^{\leq}) \setminus [e]^{\geq} = \alpha - e' \cup [e]^{\leq} \setminus [e]^{\geq}$ implying by [P1] that $\alpha - e' \cup [e]^{\leq} \approx \alpha - e' \cup [e]^{\leq} \setminus [e]^{\geq}$. Hence, $\alpha - e' \triangleright_{\approx} e$ by Definition 16.

[E4]: We need to show that $\alpha \cup [e]^{\leq} \triangleright_{\approx} e \Rightarrow \alpha \triangleright_{\approx} e$. Indeed, by Definition 16, $\alpha \cup [e]^{\leq} \triangleright_{\approx} e$ iff $(\alpha \cup [e]^{\leq}) \cup [e]^{\leq} \setminus [e]^{\geq} \approx (\alpha \cup [e]^{\leq}) \cup [e]^{\leq} \Leftrightarrow \alpha \cup [e]^{\leq} \setminus [e]^{\geq} \approx \alpha \cup [e]^{\leq}$ iff $\alpha \triangleright_{\approx} e$.

Lemma 24 Let $\mathcal{P} = (E, \leq, \approx)$ be a DPES, and let $\alpha \in \mathcal{L}_{fin}(E)$. Then $\alpha \approx$

$ST_{\triangleright \approx}(\alpha)$.

Proof By Theorem 23 and Lemma 18, $ST_{\triangleright \approx}(\alpha)$ is a configuration. Let $\alpha \setminus ST_{\triangleright \approx}(\alpha) = \{e_1, \dots, e_n\}$. By Definition 16, $\alpha \triangleright \approx e_i \Rightarrow \alpha \approx \alpha \setminus [e_i]^\geq$. Clearly, $\bigcap_{i=1}^n \alpha \setminus [e_i]^\geq \subseteq ST_{\triangleright \approx}(\alpha) \subseteq \alpha$, implying by [P1] and $\alpha \approx \alpha \setminus [e_i]^\geq$ that $\alpha \approx ST_{\triangleright \approx}(\alpha)$.

The next proposition gives a characterization of \approx via $\triangleright \approx$, needed for establishing an isomorphism between DPESs and DEESs.

Proposition 25 In a DPES $\mathcal{P} = (E, \leq, \approx)$, $\alpha \approx \beta$ iff $ST_{\triangleright \approx}(\alpha) = ST_{\triangleright \approx}(\beta)$.

Proof

(\Leftarrow) Immediate from Lemma 24.

(\Rightarrow) By Lemma 24, $\alpha \approx ST_{\triangleright \approx}(\alpha) \approx ST_{\triangleright \approx}(\beta)$. Suppose on the contrary that $ST_{\triangleright \approx}(\alpha) \neq ST_{\triangleright \approx}(\beta)$, say $\exists e \in ST_{\triangleright \approx}(\alpha) \setminus (ST_{\triangleright \approx}(\alpha) \cap ST_{\triangleright \approx}(\beta))$. By [P1], $ST_{\triangleright \approx}(\alpha) \approx ST_{\triangleright \approx}(\alpha) \cap ST_{\triangleright \approx}(\beta)$. Further, $ST_{\triangleright \approx}(\alpha) \cap ST_{\triangleright \approx}(\beta) \subseteq \alpha \setminus [e]^\geq \subseteq \alpha$ (since $ST_{\triangleright \approx}(\alpha) \cap ST_{\triangleright \approx}(\beta)$ is left-closed), hence again by [P1] $\alpha \cup [e]^\leq \setminus [e]^\geq = \alpha \setminus [e]^\geq \approx \alpha = \alpha \cup [e]^\leq$, implying by Definition 16 that $\alpha \triangleright \approx e$, contradicting $e \in ST_{\triangleright \approx}(\alpha)$.

Theorem 26 (Equivalence of DPESs and DEESs)

- (1) For any DEES $\mathcal{C} = (E, \leq, \triangleright)$, $\mathcal{C}_{P_C} = \mathcal{C}$.
- (2) For any DPES $\mathcal{P} = (E, \leq, \approx)$, $\mathcal{P}_{C_P} = \mathcal{P}$.

Proof Note that the translations of DEESs into DPES and vice versa does not effect the sets of events and the causality relation. Thus $\mathcal{C}_{P_C} = (E, \leq, \triangleright_{\approx \triangleright})$ and $\mathcal{P}_{C_P} = (E, \leq, \approx_{\triangleright \approx})$. We need to show that (1) $\triangleright = \triangleright_{\approx \triangleright}$ and (2) $\approx = \approx_{\triangleright \approx}$.

- (1) We need to prove that $\alpha \triangleright e$ in \mathcal{C} iff $\alpha \triangleright_{\approx \triangleright} e$ in \mathcal{C}_{P_C} . By [E4], [E1] and [E3], $\alpha \triangleright e$ iff $\alpha \cup [e]^\leq \triangleright e$ iff $\alpha \cup [e]^\leq \triangleright [e]^\geq$. By Proposition 21, the latter holds iff $ST_{\triangleright}(\alpha \cup [e]^\leq) = ST_{\triangleright}(\alpha \cup [e]^\leq \setminus [e]^\geq)$, which, by Definition 17 means that $\alpha \cup [e]^\leq \approx_{\triangleright} \alpha \cup [e]^\leq \setminus [e]^\geq$, meaning $\alpha \triangleright_{\approx \triangleright} e$ by Definition 16.
- (2) We need to prove that $\alpha \approx \beta$ in \mathcal{P} iff $\alpha \approx_{\triangleright \approx} \beta$ in \mathcal{P}_{C_P} . By Proposition 25 and Definition 17, $\alpha \approx \beta$ iff $ST_{\triangleright \approx}(\alpha) = ST_{\triangleright \approx}(\beta)$ iff $\alpha \approx_{\triangleright \approx} \beta$.

From now on, we will merge DEESs $\mathcal{C} = (E, \leq, \triangleright)$ and DPESs $\mathcal{P} = (E, \leq, \approx)$ into one model $(E, \leq, \triangleright, \approx)$, where (E, \leq) is a Prime Event Structure, the erasure relation \triangleright satisfies axioms [E0] - [E4], and $\approx = \approx_{\triangleright}$. The obtained results allow us to assume that the permutation-equivalence \approx satisfies axioms [P0] - [P3], and that $\triangleright = \triangleright_{\approx}$. We call such models again DEESs, and denote by \mathcal{C} .

4 Relating Conflict-Free Stable Transition and Event models

We now define translations between AZDFSs and DEESs and show that they commute, implying equivalence of the two computational models.

4.1 Translation of ASDRSs and AZDFSs into DEESs

Definition 27 With an ASDRS \mathcal{R} and its corresponding AZDFS $\mathcal{F} = \mathcal{F}_{\mathcal{R}} = (\mathcal{R}, \simeq_z, \hookrightarrow_z)$, we associate the DEES $\mathcal{C}_{\mathcal{R}} = \mathcal{C}_{\mathcal{F}} = (E_{\mathcal{F}}, \leq, \triangleright, \approx)$, where

- $E_{\mathcal{F}} = FAM(\mathcal{F})$, the set of all families of \mathcal{F} ;
- $\phi < \psi$ iff $\phi \hookrightarrow_z \psi$;
- $\alpha \approx \beta$ iff $\forall \phi \in \alpha \cup \beta \setminus \alpha \cap \beta, \exists P : t_{\emptyset} \twoheadrightarrow s : FAM(P) \subseteq \alpha \cap \beta$ and P discards a canonical element of ϕ (see Theorem 13);
- $\alpha \triangleright \phi$ iff $\exists P : t_{\emptyset} \twoheadrightarrow s : FAM(P) \subseteq \alpha$ and P discards a canonical element of ϕ .

We need to check that $\mathcal{C}_{\mathcal{F}}$ in the above definition is indeed a DEES. We need a number of technical lemmas to achieve this.

Lemma 28 Let $Q : t \twoheadrightarrow s, v \subseteq t, Q' \in STV(Q)$, and let v be Q -inessential, in an ASDRS. Then Q' discards v .

Proof Since v is Q -inessential, $v/Q = \emptyset$ and v is Q -unneded. Since Q' is Q -needed, it is external to v by Lemma 7.(4), and $v/Q' = v/Q = \emptyset$, i.e., Q' discards v .

Lemma 29 Let $Q : t \twoheadrightarrow s$ be standard and let $t \xrightarrow{u} o$, in an ASDRS. Then Q/u is standard.

Proof By induction on the length of Q . Let $Q' = Q/u$. Suppose on the contrary that Q' is not standard and let w' be the last Q' -unneded step in Q' , say $Q' = Q'_1 + w' + Q'_2$, and let $Q = Q_1 + w + Q_2$, where $w' = w/u'$ and $u' = u/Q_1$ (see the figure).

$$\begin{array}{ccccc}
 t & \xrightarrow{Q_1} & & \xrightarrow{w} & \xrightarrow{Q_2} \\
 u \downarrow & & u' \downarrow & & \downarrow \\
 o & \xrightarrow{Q'_1} & & \xrightarrow{w'} & \xrightarrow{Q'_2}
 \end{array}$$

By Lemma 7.(2), w' is not the last step of Q' . So let $Q_2 = v + Q_3$ and $Q'_2 = v' + Q'_3$. By the choice of w' , v' is Q' -needed, and w' does not create v' by Lemma 7.(3). Since $w + Q_2$ is self-needed and $w' + Q'_2$ is not, $u' \neq \emptyset$. Hence,

by [stability], w does not create v . Thus w' and v' , as well as w and v , can be permuted in $w' + v' + Q'_3$ and $w + v + Q_3$, respectively, yielding $v'^* + w'^* + Q'_3$ and $v^* + w^* + Q_3$ (see the figure below). Then by Lemma 7.(4), $w^* + Q_3$ is self-needed and $w'^* + Q'_3$ is not self-needed (and therefore $u'' = u'/v^* \neq \emptyset$), contradicting the induction assumption.

$$\begin{array}{ccccccc}
& & Q_1 & \xrightarrow{v^*} & & \xrightarrow{w^*} & Q_3 \\
t & \xrightarrow{\quad} & & & & & \\
u \downarrow & & u' \downarrow & & u'' \downarrow & & \downarrow \\
o & \xrightarrow{\quad} & & \xrightarrow{v'^*} & & \xrightarrow{w'^*} & Q'_3 \\
& & Q'_1 & & & &
\end{array}$$

Lemma 30 Let $Q : t \rightarrow s$, $P : t \rightarrow o$, $v \subseteq o$, $Q' \in STV(Q)$, and let Pv be Q -inessential, in an ASDRS. Then Q' discards Pv .

Proof By Lemma 29, $Q'/P \in STV(Q/P)$. And since v is Q/P -inessential, Q'/P discards v by Lemma 28.

Lemma 31 Let $P : t \rightarrow s$, $Q : t \xrightarrow{Q'} o \xrightarrow{u} e \xrightarrow{v} s'$, let $Q'u$ be P -inessential, and let u create v , in an ASDRS. Then $(Q' + u)v$ is P -inessential too.

Proof Let $P' = ST(P)$, $P'' = P'/Q'$, and $P^* = P''/u$. By Lemma 30, P'' discards u . So let w be the step of P'' that discards a residual u' of u , and let v' be the corresponding residual of v along P^* , see the figure. If $v' \neq \emptyset$, then $v'/(w/u') = \emptyset$ by [stability] (since $u'/w = \emptyset$). So, in any case, $v/P^* = \emptyset$. By the Stability Lemma, P^* is external to v (since P'' is external to u). Hence P^* discards v , i.e. v is P -inessential.

$$\begin{array}{ccccccc}
& & & \xrightarrow{P'} & & & \\
Q' \downarrow & & & & & & \downarrow \\
& \xrightarrow{\quad} & w \in P'' & \xrightarrow{\quad} & & & \\
u \downarrow & & u' \downarrow & & \emptyset \downarrow & & \emptyset \downarrow \\
& \xrightarrow{\quad} & w' \in P^* & \xrightarrow{\quad} & & & \\
v \downarrow & & v' \downarrow & & \emptyset \downarrow & & \emptyset \downarrow \\
& \xrightarrow{\quad} & & \xrightarrow{\quad} & & &
\end{array}$$

Lemma 32 Let $P : t_\emptyset \rightarrow s$, ϕ be a family in an AZDFS \mathcal{F} , let $Qv, Q'v' \in \phi$, and let Qv be canonical. If Qv is P -inessential, then so is $Q'v'$. Further, if $Q' \trianglelefteq P$, then the converse is also true.

Proof By the Affine Extraction Theorem, there is Q'' such that $Q' \approx Q + Q''$ and $v' = v/Q''$. If Qv is P -inessential, then $ST(P)/Q$ discards v by Lemma 30, and $ST(P)/Q'$ discards v' by the Cube Lemma. Conversely, if $Q' \trianglelefteq P$, then $Q''/(P/Q) = \emptyset$, and $v'/(P/Q') = v/(P/Q) = \emptyset$ again by the Cube Lemma, thus Qv is P -inessential.

Lemma 33 Let $t_\emptyset \xrightarrow{Q} s \xrightarrow{u} s'$, and $t_\emptyset \xrightarrow{P} o \xrightarrow{v} o'$, and $Qu \simeq_z Pv$, in an AZDFS \mathcal{F} . Then $v/((Q+u)/P) = \emptyset$.

Proof If $v' = v/(Q/P) = \emptyset$, then we are done. Otherwise, by the Affine Extraction Theorem, there are $Nw \simeq_z Qu$ and reductions N' and N'' such that $Q \approx N+N'$, $P \approx N+N''$, $u = w/N'$ and $v = w/N''$. Further, $N'+P/Q \approx N''+Q/P$, and by the Cube Lemma, $v' = w/(N'+P/Q)$. But u is the only N' -residual of w , thus $v' = u/(P/Q)$, implying $v/((Q+u)/P) = \emptyset$.

Lemma 34 Let $P \in STA$, $FAM(P) = F_1 \cup F_2$, $F_1 \cap F_2 = \emptyset$ and $\forall \phi \in F_2, \forall \psi \in F_1 : \phi \not\leftrightarrow \psi$, in an AZDFS \mathcal{F} . Then $\exists Q \in STV(P) : Q = Q_1 + Q_2 \wedge FAM(Q_i) = F_i, i = 1, 2$.

Proof Let u be the latest step in P such that $\phi = Fam(u) \in F_2$ and it has a next step v with $\psi = Fam(v) \in F_1$. Then $\phi \not\leftrightarrow \psi$ by the hypothesis, and u does not create v by [creation]. Hence u and v can be permuted, yielding another standard reduction by Lemma 7.(4). The resulting reduction can be transformed similarly until all steps in F_2 are pushed to the end. The transformation terminates as it does not change the length of the reduction.

Theorem 35 For any AZDFS \mathcal{F} , \mathcal{C}_F is a DEES.

Proof We need to show that $\mathcal{C}_F = (E_{\mathcal{F}}, \xleftrightarrow{\quad}, \triangleright, \approx)$ satisfies the erasure axioms [E0] – [E4], and that $\approx = \approx_{\triangleright}$.

[E0]: Immediate from Definitions 6 and 27.

[E1]: Again from Definitions 6 and 27.

[E2]: Let $\alpha \triangleright \phi$, $\alpha \triangleright \phi'$, and let $\alpha - \phi \in \mathcal{L}_{fin}(E)$. Then there are P and P' such that $FAM(P), FAM(P') \subseteq \alpha$, a canonical element Qv of ϕ is P -inessential, and a canonical element $Q'v'$ of ϕ' is P' -inessential. Let $P^* = P \sqcup P'$. Then $FAM(P^*) \subseteq \alpha$ and both Qv and $Q'v'$ are P^* -inessential. By Lemma 32, all members of ϕ and ϕ' are P^* -inessential. Since $ST(P^*)$ is P^* -essential, it does not contract members of ϕ and ϕ' . Hence $FAM(ST(P^*)) \subseteq \alpha - \phi'$, and $\alpha - \phi' \triangleright \phi$ by Definition 27.

[E3]: Let $\alpha \triangleright \phi$ and $\phi \hookrightarrow \phi'$. It is enough to consider the case when $\exists \phi'' : \phi \hookrightarrow \phi'' \hookrightarrow \phi'$. By Lemma 34 and the Affine Extraction Theorem, there is a canonical $P'v' \in \phi'$ such that $P' = P + v$, v creates v' , and $Pv \in \phi$. Further, by Definition 27, $\exists Q : FAM(Q) \subseteq \alpha$ which discards a canonical element of ϕ . Hence, by Lemmas 32, 31 and 30, $ST(Q)$ discards $P'v'$, i.e., $\alpha \triangleright \phi'$.

[E4]: Let $\alpha \cup [\phi]^< \triangleright \phi$. By Definition 27, there is a Q such that $FAM(Q) \subseteq \alpha \cup [\phi]^<$ and Q discards a canonical element Pv of ϕ . By Proposition 11 and Lemma 30, Q can be chosen standard. Let $F_2 = FAM(Q) \cap ([\phi]^< \setminus \alpha)$ and $F_1 = FAM(Q) \setminus F_2$. Since $\alpha \in \mathcal{L}_{fin}(E)$, $\forall \phi \in F_2, \psi \in F_1 : \phi \not\leftrightarrow \psi$. Hence, by Lemma 34, $\exists Q' \in STV(Q) : Q' = Q_1 + Q_2 \wedge FAM(Q_i) = F_i$. By Lemma 30, Q' discards Pv , and by Lemma 33, $Q'/P = Q_1/P$, since

$[\phi]^< = FAM(P)$. Hence Q_1 discards Pv and $FAM(Q_1) \subseteq \alpha$, i.e., $\alpha \triangleright \phi$.

It remains to show that $\approx = \approx_{\triangleright}$. Indeed, by Definition 27, $\alpha \approx \beta$ iff $\alpha \cap \beta \triangleright \alpha \cup \beta \setminus \alpha \cap \beta$, and by Proposition 21, the latter holds iff $\alpha \approx_{\triangleright} \beta$.

4.2 Translation of DEESs into ASDRSs and AZDFSs

The following translation of DEESs into ASDRSs uses a techniques resembling the representation of events in a PES as prime intervals of its configuration domain, used for construction of PESs from stable families of configurations in [37].

Definition 36 With a DEES $\mathcal{C} = (E, \leq, \triangleright, \approx)$ we associate an ASDRS $\mathcal{R}_{\mathcal{C}}$ as follows:

- The terms of $\mathcal{R}_{\mathcal{C}}$ are Lévy-equivalence classes $\langle \alpha \rangle_L, \langle \beta \rangle_L, \dots$ of configurations of \mathcal{C} ;
- The reduction relation of $\mathcal{R}_{\mathcal{C}}$ consists of sets of pairs $u = (\langle \alpha \rangle_L, e)$, where $\alpha, \alpha + e \in \mathcal{L}_{fin}(E)$ and $\alpha \not\approx \alpha + e$. Pairs $u = (\langle \alpha \rangle_L, e)$ such that $\alpha \approx \alpha + e$ are identified with the empty redex $\emptyset = (\langle \alpha \rangle_L, \emptyset)$ in $\langle \alpha \rangle_L$. A non-empty redex $(\langle \alpha \rangle_L, e)$ will also be called an *e-redex*.
- The residual relation is defined as follows: if $u = (\langle \alpha \rangle_L, e)$ and $v = (\langle \alpha \rangle_L, e')$, then $u/v = (\langle \alpha + e' \rangle_L, e)$. (Thus $u/v = \emptyset$ iff $\alpha + e' \approx \alpha + e' + e$.)

Next we show that the definition above is correct.

Theorem 37 Let $\mathcal{C} = (E, \leq, \triangleright, \approx)$ be a DEES. Then $\mathcal{R}_{\mathcal{C}}$ is an ASDRS.

Proof For any step $v : \langle \alpha \rangle_L \xrightarrow{(\langle \alpha \rangle_L, e)} \langle \alpha + e \rangle_L$ in $R_{\mathcal{C}}$, and any redex $u = (\langle \alpha \rangle_L, e')$ in $\langle \alpha \rangle_L$, u/v is either a (non-empty) redex in $\langle \alpha + e \rangle_L$, or $u/v = \emptyset$. Also, for any other redex $w = (\langle \alpha \rangle_L, e'')$ in $\langle \alpha \rangle_L$ such that $w/v \neq \emptyset$, $w/v = (\langle \alpha + e \rangle_L, e'') \neq (\langle \alpha + e \rangle_L, e') = u/v$, i.e., no redex in $\langle \alpha + e \rangle_L$ is a residual of more than one redex in $\langle \alpha \rangle_L$. Further, $v/v = (\langle \alpha + e \rangle_L, e) = \emptyset$. So to show that $R_{\mathcal{C}}$ is an ASDRS, it remains to show [FD], [weak acyclicity], and [stability].

[FD] let U be a set of redexes in $\langle \alpha \rangle_L$, let e_0, \dots, e_{l-1} be an enumeration of U , and let $\alpha_n = \alpha + e_0 + \dots + e_{n-1}$ ($0 \leq n \leq l$). Then a complete development of U has a form $P : \langle \alpha_0 \rangle_L \xrightarrow{(\langle \alpha_0 \rangle_L, e_0)} \langle \alpha_1 \rangle_L \xrightarrow{(\langle \alpha_1 \rangle_L, e_1)} \dots \rightarrow \langle \alpha_k \rangle_L$, where for any $m = k, \dots, l-1$, $(\langle \alpha_0 \rangle_L, e_m)/P = \emptyset$, that is, $\langle \alpha_k \rangle_L = \langle \alpha_k + e_m \rangle_L$. We show by induction on $j = l - k$ that $\alpha_k \approx \alpha_k + e_k + \dots + e_{l-1}$. The case $j = 1$ (i.e., $\alpha_k \approx \alpha_k + e_k$) is immediate. By the induction assumption, $\alpha_k \approx \alpha_{l-2} \approx \alpha_{l-2} + e_{l-2} \approx \alpha_{l-2} + e_{l-1}$. By [P2], $\alpha_{l-2} + e_{l-2} + e_{l-1} \approx \alpha_{l-2} + e_{l-1}$. Hence, $\alpha_k \approx \alpha_{l-2} + e_{l-2} + e_{l-1} = \alpha_l$. Thus every complete development of U

ends at $\langle \alpha_k \rangle_L = \langle \alpha_l \rangle_L$. Further, for any $u = (\langle \alpha \rangle_L, e)$, $u/P = (\langle \alpha_k \rangle_L, e)$, i.e., u/U does not depend on the particular complete development of U , hence [FD] holds.

[weak acyclicity] Let $u = (\langle \alpha \rangle_L, e)$, $v = (\langle \alpha \rangle_L, e')$, $e \neq e'$, and suppose $u/v = v/u = \emptyset$. That is, $\alpha + e + e' \approx \alpha + e \approx \alpha + e'$. Then, by [P1], $\alpha + e \approx \alpha + e' \approx \alpha$, i.e., $u = v = \emptyset$.

[stability] Suppose on the contrary that \mathcal{R}_C is not stable. Then there are $u \neq v$, w , w' and w'' such that w and w' are created by u and v respectively, and $w'' = w/v' = w'/u' \neq \emptyset$, where $u' = u/v$ and $v' = v/u$. That is, there are $u = (\langle \alpha \rangle_L, e)$, $v = (\langle \alpha \rangle_L, e')$, $w = (\langle \alpha + e \rangle_L, e'')$, $w' = (\langle \alpha + e' \rangle_L, e'')$, and $w'' = (\langle \alpha + e + e' \rangle_L, e'')$ such that $\langle \alpha \rangle_L$, $\langle \alpha + e \rangle_L$ and $\langle \alpha + e + e'' \rangle_L$ are different; so are $\langle \alpha \rangle_L$, $\langle \alpha + e' \rangle_L$ and $\langle \alpha + e' + e'' \rangle_L$; and $\langle \alpha + e + e' \rangle_L \neq \langle \alpha + e + e' + e'' \rangle_L$ (see the diagram). Further, $(\langle \alpha \rangle_L, e'')$ is not a redex since w is a created redex. Hence either $\alpha + e'' \notin \mathcal{L}_{fin}(\mathcal{C})$, or $\alpha + e'' \in \mathcal{L}_{fin}(\mathcal{C})$ and $\alpha \approx \alpha + e''$; but the latter would imply by [P2] that $\alpha + e + e'' \approx \alpha + e$, contradicting the fact that w is a non-empty redex. Thus $\alpha + e + e''$ and $\alpha + e' + e''$ are both configurations, but their intersection $\alpha + e''$ is not – a contradiction, since the intersection of left-closed subsets remains left-closed.

$$\begin{array}{ccccc}
 \langle \alpha \rangle_L & \xrightarrow{u} & \langle \alpha + e \rangle_L & \xrightarrow{w} & \langle \alpha + e + e'' \rangle_L \\
 v \downarrow & & \downarrow v' & & \downarrow \\
 \langle \alpha + e' \rangle_L & \xrightarrow{u'} & \langle \alpha + e + e' \rangle_L & \xrightarrow{w''} & \langle \alpha + e + e' + e'' \rangle_L \\
 w' \downarrow & & & & \downarrow \\
 \langle \alpha + e' + e'' \rangle_L & \xrightarrow{\quad\quad\quad} & & & \langle \alpha + e + e' + e'' \rangle_L
 \end{array}$$

The theorem is now proved.

Clearly, an initial reduction in an ASDRS \mathcal{R}_C can be represented as a left-closed enumeration e_1, \dots, e_k of a configuration α where no event e_i in the sequence is *vacuous*, i.e., $\forall i : \{e_1, \dots, e_{i-1}\} \not\triangleright e_i$; the event e_i represents the *non-empty* redex $(\{e_1, \dots, e_{i-1}\}, e_i)$. We call such a sequence also an *initial reduction* or an α -*reduction* in the DEES \mathcal{C} , written $[\alpha]$. Thus we have the following lemma:

Lemma 38 Let \mathcal{C} be a DEES, and let \mathcal{R}_C be its corresponding ASDRS. Then there is a 1 – 1 correspondence between initial reductions in \mathcal{C} and \mathcal{R}_C .

Note that, unlike PESs, not every configuration in a DEES has a reduction. For example, consider the DEES consisting of four events $\{e_1, e_2, e'_1, e'_2\}$, ordered as follows: $e_1 < e'_1$ and $e_2 < e'_2$, and let \triangleright be given by: $\alpha \triangleright e'_1$ iff $e_2 \in \alpha$, and $\alpha \triangleright e'_2$ iff $e_1 \in \alpha$. Then $\{e_1, e_2, e'_1, e'_2\}$ is a configuration, but any of its left-closed enumerations, such as e_1, e_2, e'_1, e'_2 , ends either with e'_1 or with e'_2 , and we have $\{e_1, e_2, e'_2\} \triangleright e'_1$ and $\{e_1, e_2, e'_1\} \triangleright e'_2$. This is not surprising as the vacuous events in left-closed sequences correspond to empty reductions in the

corresponding ASDRS, performed in the ‘garbage’ (erased redexes). However, we have the following lemma:

Lemma 39 Let $\alpha \in \mathcal{L}_{fin}(E)$, in a DEES. Then any left-closed enumeration of $ST(\alpha)$ is a reduction.

Proof Let e_0, \dots, e_n be a left-closed enumeration of $ST(\alpha)$. If there was an i such that $e_0, \dots, e_{i-1} \triangleright e_i$, then $\alpha \triangleright e_i$ by [E1], contradicting $e_i \in ST(\alpha)$.

Definition 40 With a DEES $\mathcal{C} = (E, \leq, \triangleright, \approx)$, we associate a DFS $\mathcal{F}_{\mathcal{C}} = (\mathcal{R}_{\mathcal{C}}, \simeq_{\mathcal{C}}, \hookrightarrow_{\mathcal{C}})$ as follows:

- $\mathcal{R}_{\mathcal{C}}$ is the DRSs defined in Definition 36.
- $\forall e \in E$, we define the $\simeq_{\mathcal{C}}$ -family ϕ_e corresponding to e , or the *e-family*, as the set of all e -redexes $([\alpha], e)$. That is, $([\alpha], e) \simeq_{\mathcal{C}} ([\beta], e')$ iff $e = e'$.
- Define $\hookrightarrow_{\mathcal{C}}$ by: $\phi_e \hookrightarrow_{\mathcal{C}} \phi_{e'}$ iff $e < e'$.

Theorem 41 Let $\mathcal{C} = (E, \leq, \triangleright, \approx)$ be a DEES, let $\mathcal{F}_{\mathcal{C}} = (\mathcal{R}_{\mathcal{C}}, \simeq_{\mathcal{C}}, \hookrightarrow_{\mathcal{C}})$, and let $\mathcal{F}_{\mathcal{R}_{\mathcal{C}}} = (\mathcal{R}_{\mathcal{C}}, \simeq_z, \hookrightarrow_z)$. Then $\mathcal{F}_{\mathcal{C}} = \mathcal{F}_{\mathcal{R}_{\mathcal{C}}}$, hence $\mathcal{F}_{\mathcal{C}}$ is an AZDFS.

Proof We need to show that (a) $\simeq_z = \simeq_{\mathcal{C}}$ and (b) $\hookrightarrow_z = \hookrightarrow_{\mathcal{C}}$.

- (a) By Definition 36, $([\alpha], e) \simeq_z ([\beta], e') \Rightarrow e = e'$, implying $\simeq_z \subseteq \simeq_{\mathcal{C}}$. For the converse, consider two redexes $([\alpha], e) \simeq_{\mathcal{C}} ([\beta], e)$. Then $([\alpha \cap \beta], e)$ is also a redex since $\alpha \cap \beta + e = \alpha + e \cap \beta + e \in \mathcal{L}_{fin}(E)$, and $([\alpha], e)$ and $([\beta], e)$ are its residuals. (Note that $(\langle \alpha \cap \beta \rangle_L, e) \neq \emptyset$ since $(\langle \alpha \cap \beta \rangle_L, e) = \emptyset \Rightarrow (\langle \alpha \rangle_L, e) = \emptyset$ by [P2].) Hence $([\alpha], e) \simeq_z ([\beta], e)$.
- (b) In order to show that $\phi_e \hookrightarrow_{\mathcal{C}} \phi_{e'} \Leftrightarrow \phi_e \hookrightarrow_z \phi_{e'}$, we need to show that $e < e'$ iff $e \in [\gamma]$ for any $([\gamma], e') \in \phi_{e'}$ (by (a), $\phi_e, \phi_{e'}$ are families both w.r.t. $\simeq_{\mathcal{C}}$ and \simeq_z). Sufficiency is immediate since $\gamma + e'$ is left closed. For the converse, we have from Theorem 13 that $([[e']^<], e')$ is a canonical element of $\phi_{e'}$ (when $\phi_{e'}$ is considered as a \simeq_z -family), hence $e < e'$.

4.3 Equivalence of the transition and event models

We now establish an equivalence between DEESs and AZDFSs, implying together with the results of previous sections equivalence of the considered four transition and event models – ASDRSs, AZDFSs, DEESs and DPESs.

Let \mathcal{R} be a DRS. Let us define the *reduction tree* of \mathcal{R} to be a tree whose nodes are labelled with terms in \mathcal{R} , whose root is labelled with t_{\emptyset} , the initial term of \mathcal{R} , and whose arcs correspond to redexes in \mathcal{R} . Thus sons of a node are target terms of redexes whose source term is the label of that node. (Reduction trees are considered up to the order of arcs coming out of a node: the order

in which the arcs are drawn is irrelevant.) Note that different nodes can be labelled with the same term, so even if \mathcal{R} has cycles, its reduction tree is a tree. Thus the branches of the reduction tree of \mathcal{R} are in 1 – 1 correspondence with initial reductions in \mathcal{R} . We call two such trees *isomorphic* if the trees obtained by removing all labels are identical. Hence, we can define two DRSS to be *isomorphic* if there is an isomorphism between their reduction trees preserving the residual relation. An isomorphism between two DFSSs must further preserve the family and contribution relations.

Lemma 42 Let \mathcal{F} be an AZDFS and let $\mathcal{C}_{\mathcal{F}}$ be its corresponding DEES.

- (1) For any initial reduction $[\alpha] = e_0, e_1, \dots, e_n$ in $\mathcal{C}_{\mathcal{F}}$ there is a unique initial reduction $g([\alpha]) = P : t_\emptyset = t_0 \xrightarrow{u_0} t_1 \xrightarrow{u_1} \dots \xrightarrow{u_{n-1}} t_n$ in \mathcal{F} such that $Fam(P_i u_i) = e_i$, where $P_i : t_\emptyset = t_0 \xrightarrow{u_0} t_1 \xrightarrow{u_1} \dots \xrightarrow{u_{i-1}} t_i$. (Thus $FAM(P) = \alpha$.)
- (2) Let $P : t_\emptyset = t_0 \xrightarrow{u_0} t_1 \xrightarrow{u_1} \dots \xrightarrow{u_{n-1}} t_n$ in \mathcal{F} , and let $e_i = \phi_i = Fam(P_i u_i)$, where $P_i : t_\emptyset = t_0 \xrightarrow{u_0} t_1 \xrightarrow{u_1} \dots \xrightarrow{u_{i-1}} t_i$. Then $f(P) = e_0, e_1, \dots, e_n$ is a reduction in $\mathcal{C}_{\mathcal{F}}$.
- (3) the translation functions g and f between reductions in \mathcal{F} and $\mathcal{C}_{\mathcal{F}}$ commute: for any initial reduction P in \mathcal{F} , $g(f(P)) = P$, and for any reduction $[\alpha]$ in $\mathcal{C}_{\mathcal{F}}$, $f(g([\alpha])) = [\alpha]$.

Proof

- (1) By definition of a reduction in a DEES, e_0, e_1, \dots, e_n is left-closed, and $\{e_0, \dots, e_{j-1}\} \not\triangleright e_j$ for any $j = 0, \dots, n$. Take for u_0 the redex in t_\emptyset such that $Fam(\emptyset u_0) = e_0$; existence of such a redex is immediate from the definition of $\mathcal{C}_{\mathcal{F}}$, and uniqueness follows from [initial]. Suppose u_1, \dots, u_{i-1} are already defined so that $Fam(P_k u_k) = e_k$ for all $1 \leq k < i$. We need to show that $\exists u_i \in t_i$ such that $Fam(P_i u_i) = e_i$; uniqueness of such a redex follows from the separability of \mathcal{F} (see Theorem 12) and from the fact that in separable AZDFSs, a term may contain at most one redex of a given family (see Lemma 6.1 in [18], or Lemma 5.2 in [20]). Let $N_i v_i$ be a canonical element of e_i which exists by the Affine Extraction Theorem. By the same theorem, N_i contracts exactly one redex in every contributor family of e_i (and does not contract any other redex). Thus, by [contribution], $FAM(N_i) \subseteq FAM(P_i)$, and $N_i \trianglelefteq P_i$ by Proposition 11. Now we can take $u_i = v_i / (P_i / N_i)$. If on the contrary $u_i = \emptyset$, then P_i would discard $N_i v_i$ (since P_i / N_i is external to v_i as $Fam(N_i v_i) \notin FAM(P_i)$), and we would have by Definition 27 that $\{e_0, \dots, e_{i-1}\} \triangleright e_i$, contradicting the assumption. Thus a reduction P as above exists, and is unique.
- (2) We need to show that $f(P) = e_0, e_1, \dots, e_n$ is left-closed and does not contain vacuous events (by Definition 27, $\phi_i = e_i$ are events in $\mathcal{C}_{\mathcal{F}}$). By [contribution], P_i contracts at least one redex in every contributor family of ϕ_i , hence $f(P)$ is left-closed. If on the contrary say ϕ_j is vacuous, i.e., $\{\phi_0, \dots, \phi_{j-1}\} \triangleright \phi_j$, then by Definition 27 there is an initial reduction

Q such that $FAM(Q) \subseteq \{\phi_0, \dots, \phi_{j-1}\}$ and Q discards a canonical element $N_j v_j$ of ϕ_j . By Proposition 11, $N_j, Q \trianglelefteq P_j$, and by Definition 6, P_j erases $N_j v_j$, i.e., $v_j/(P_j/N_j) = \emptyset$. Hence $u_j/((N_j + v_j)/P_j) = u_j/\emptyset = u_j$, contradicting Lemma 33 (since $N_j v_j \simeq_z P_j u_j$).

(3) Immediate from the constructions in (1) and (2).

Theorem 43 (Equivalence of AZDFSs and DEESs)

- (1) For any DEES \mathcal{C} , \mathcal{C} and $\mathcal{C}_{\mathcal{F}_{\mathcal{C}}}$ are isomorphic.
- (2) For any AZDFS \mathcal{F} , \mathcal{F} and $\mathcal{F}_{\mathcal{C}_{\mathcal{F}}}$ are isomorphic.

Proof

- (1) Let $\mathcal{C} = (E, \leq, \triangleright, \approx)$ and $\mathcal{C}' = \mathcal{C}_{\mathcal{F}_{\mathcal{C}}} = (E', \leq', \triangleright', \approx')$. By Definitions 40 and 27, $E' = \{\phi_e | e \in E\}$, where ϕ_e is the e -family of $\mathcal{F}_{\mathcal{C}}$. We show that $f : e \mapsto \phi_e$ is an isomorphism between \mathcal{C} and \mathcal{C}' . Since f clearly an isomorphism between E and E' , and since \mathcal{C}' is a DEES by Theorems 35 and 41, we need to show that (a) $e \leq e'$ iff $\phi_e \leq' \phi_{e'}$; and (b) $\alpha \approx \beta$ iff $\phi_\alpha \approx' \phi_\beta$, where $\phi_\alpha = \{\phi_e | e \in \alpha\}$ and $\phi_\beta = \{\phi_e | e \in \beta\}$.
 - (a) Immediate from Definitions 40 and 27.
 - (b) By Definition 27, $\phi_\alpha \approx' \phi_\beta$ iff $\forall e \in \alpha \cup \beta \setminus \alpha \cap \beta$, there is a reduction $[\gamma]$ in $\mathcal{F}_{\mathcal{C}}$ such that $FAM([\gamma]) \subseteq \phi_\alpha \cap \phi_\beta$ and $[\gamma]$ discards a canonical element $[[e]^{<}e]$ of ϕ_e (see Theorem 13). By Definitions 6 and 36, the latter means that $e \notin \gamma$, $\gamma \subseteq \alpha \cap \beta$, and $\gamma \cup [e]^{<} \approx \gamma \cup [e]^{<}$, thus by Definition 16, and Lemmas 39 and 19 we need to prove that $\alpha \approx \beta$ iff \exists *standard* $\gamma : \gamma \subseteq \alpha \cap \beta \wedge \gamma \triangleright \alpha \cup \beta \setminus \alpha \cap \beta$. But this is immediate from Proposition 21 (since $\approx = \approx_{\triangleright}$) and Lemma 19.
- (2) By Theorem 41 and Theorem 35, $\mathcal{F}_{\mathcal{C}_{\mathcal{F}}}$ is an AZDFS, hence it is enough to show that the underlying ASDRSs \mathcal{R} and $\mathcal{R}_{\mathcal{C}_{\mathcal{F}}}$ of \mathcal{F} and $\mathcal{F}_{\mathcal{C}_{\mathcal{F}}}$ are isomorphic. This follows immediately from Lemmas 42 and 38.

5 Lévy's Configuration Domains

Next we define Lévy's configuration domains in DEESs and show that they are isomorphic to Lévy's reduction spaces in corresponding ASDRSs.

- Definition 44** (1) Let \mathcal{C} be a DEES. *Lévy's configuration domain* $\mathcal{L}_{fin}^{\triangleleft}(\mathcal{C}) = (\mathcal{L}_{fin}^{\approx}(\mathcal{C}), \triangleleft)$, where $\mathcal{L}_{fin}^{\approx}(\mathcal{C}) = \mathcal{L}_{fin}(\mathcal{C})/\approx = \{\langle \alpha \rangle_L | \alpha \in \mathcal{L}_{fin}(\mathcal{C})\}$ and \triangleleft is a partial order defined by: $\langle \alpha \rangle_L \triangleleft \langle \beta \rangle_L$ iff $\exists \beta' \in \langle \beta \rangle_L : \alpha \subseteq \beta'$.
- (2) Let \mathcal{R} be an ASDRS, and let $\mathcal{L}_{fin}(\mathcal{R})$ be the set of initial reductions in \mathcal{R} . *Lévy's reduction space* $\mathcal{L}_{fin}^{\triangleleft}(\mathcal{R}) = (\mathcal{L}_{fin}^{\approx}(\mathcal{R}), \triangleleft)$, where $\mathcal{L}_{fin}^{\approx}(\mathcal{R}) = \mathcal{L}_{fin}(\mathcal{R})/\approx = \{\langle P \rangle_L | P \in \mathcal{L}_{fin}(\mathcal{R})\}$, and $\langle P \rangle_L \triangleleft \langle Q \rangle_L$ iff $P \trianglelefteq Q$.

Note that if $\langle \alpha \rangle_L \sqsubseteq \langle \beta \rangle_L$, $\alpha \approx \alpha'$ and $\beta \approx \beta'$, then $\langle \alpha' \rangle_L \sqsubseteq \langle \beta' \rangle_L$, implying that Definition 44.(1) is correct: Indeed, $\langle \alpha \rangle_L \sqsubseteq \langle \beta \rangle_L$ iff $\exists \beta'' \approx \beta : \alpha \subseteq \beta''$, and by Proposition 21 and [E1], $\alpha \approx \alpha' \Rightarrow \alpha \triangleright (\alpha' \setminus \alpha) \Rightarrow \beta'' \triangleright (\alpha' \setminus \alpha) \Rightarrow \beta' \approx \beta \approx \beta'' \approx \beta'' \cup (\alpha' \setminus \alpha) = \beta'' \cup \alpha'$, thus $\langle \alpha' \rangle_L \sqsubseteq \langle \beta' \rangle_L$.

Lemma 45 Let P be an initial reduction in an AZDFS \mathcal{F} . Then, in the corresponding DEES $\mathcal{C}_{\mathcal{F}}$, $FAM(ST(P)) = ST(FAM(P))$.

Proof First note that, by [contribution] and the definition of \leq (Definition 27), for any initial reduction Q , $FAM(Q)$ is indeed a configuration.

- (\subseteq) Let $\phi \in FAM(ST(P))$. By Proposition 11, $\phi \in SFAM(P)$, i.e., there is $P'v' \in \phi$ that is contracted in P and is P -needed, thus is P -essential. Let P^*v^* be a canonical element of ϕ . Then it is P -essential by Lemma 32. For any initial reduction Q with $FAM(Q) \subseteq FAM(P)$, $Q \leq P$ by Proposition 11, thus P^*v^* is Q -essential too. Hence Q does not discard P^*v^* , implying by Definition 27 that $FAM(P) \not\triangleright \phi$, i.e., $\phi \in ST(FAM(P))$.
- (\supseteq) Let $\phi \in ST(FAM(P))$. Then $\phi \in FAM(P)$ and $FAM(P) \not\triangleright \phi$, which by Definition 27 means that there is no initial Q with $FAM(Q) \subseteq FAM(P)$ such that Q discards a canonical element P^*v^* of ϕ . In particular, $ST(P)$ does not discard P^*v^* , and by Lemmas 32 and 7.(1), the redex $P'v' \in \phi$ contracted in P is P -needed, i.e., $\phi \in SFAM(P)$; thus $\phi \in FAM(ST(P))$ by Proposition 11.

Lemma 46 Let Q, Q' be initial reductions in an AZDFS \mathcal{F} . Then $Q \approx Q'$ in \mathcal{F} iff $FAM(Q) \approx FAM(Q')$ in the corresponding DEES $\mathcal{C}_{\mathcal{F}} = (E_{\mathcal{F}}, \leq, \triangleright, \approx)$.

Proof

- (\Rightarrow) By Definition 27, we need to show that $\forall \phi \in FAM(Q) \cup FAM(Q') \setminus FAM(Q) \cap FAM(Q')$, an initial reduction P with $FAM(P) \subseteq FAM(Q) \cap FAM(Q')$ discards a canonical element Nv of ϕ . Take $P = ST(Q)$. By Proposition 11, $FAM(P) \subseteq FAM(Q) \cap FAM(Q')$. Let Q^*v^* be an element of ϕ contracted in Q . Since by Proposition 11 $\phi \notin FAM(P) = SFAM(Q)$, Q^*v^* is Q -unneeded, hence Q -inessential by Lemma 7.(1). Hence, by Lemma 30, P discards Q^*v^* . By the Affine Extraction Theorem, $\exists N' : N + N' \approx Q^*$ and $v^* = v/N'$. Since $Q^* \leq Q \approx P$, $N'/(P/N) = \emptyset$, hence $v/(P/N) = v^*/(P/Q^*) = \emptyset$ by the Cube Lemma. Since $\phi \notin FAM(P)$, P/N is external to v , thus P discards Nv .
- (\Leftarrow) By Proposition 25, $FAM(Q) \approx FAM(Q')$ implies that $ST(FAM(Q)) = ST(FAM(Q'))$; hence, by Lemma 45, $FAM(ST(Q)) = FAM(ST(Q'))$, implying by Proposition 11 that $Q \approx Q'$.

Lemma 47 Let \mathcal{F} be an AZDFS, let e_0, e_1, \dots, e_n be a left-closed enumeration of $\alpha \in \mathcal{L}_{fin}(\mathcal{C}_{\mathcal{F}})$, and let e_{i_0}, \dots, e_{i_m} be obtained from e_0, e_1, \dots, e_n by eliminating all vacuous events, i.e., $e_j \in \{e_{i_0}, \dots, e_{i_m}\}$ iff $\{e_0, \dots, e_{j-1}\} \not\triangleright e_j$.

Then e_{i_0}, \dots, e_{i_m} is a reduction in $\mathcal{C}_{\mathcal{F}}$ and $\{e_{i_0}, \dots, e_{i_m}\} \approx \{e_0, \dots, e_n\}$.

Proof If there were l, k such that $e_l < e_{i_k}$ (hence $l < i_k$) and $l \neq i_0, \dots, i_{k-1}$, then $\{e_0, \dots, e_{l-1}\} \triangleright e_l$, $\{e_0, \dots, e_{i_k-1}\} \not\triangleright e_{i_k}$, and by [E1], $\{e_0, \dots, e_{i_k-1}\} \triangleright e_l$, contradicting [E3]. Hence e_{i_0}, \dots, e_{i_m} is left-closed. Since $\{e_0, \dots, e_{i_k-1}\} \not\triangleright e_{i_k}$, $\{e_{i_0}, \dots, e_{i_{k-1}}\} \not\triangleright e_{i_k}$ by [E1], thus e_{i_0}, \dots, e_{i_m} is a reduction in $\mathcal{C}_{\mathcal{F}}$. Finally, $\{e_{i_0}, \dots, e_{i_m}\} \approx \{e_0, \dots, e_n\}$ follows from Proposition 21.

Proposition 48 Let P, Q be initial reductions in an ASDRS \mathcal{R} . Then $\langle P \rangle_L \trianglelefteq \langle Q \rangle_L$ iff $\langle FAM(P) \rangle_L \trianglelefteq \langle FAM(Q) \rangle_L$ in $\mathcal{C}_{\mathcal{R}}$.

Proof

- (\Rightarrow) By Proposition 11, $\langle P \rangle_L \trianglelefteq \langle Q \rangle_L$ iff $\exists Q' \approx Q : FAM(P) \subseteq FAM(Q')$, which by Definition 44 holds iff $\langle FAM(P) \rangle_L \trianglelefteq \langle FAM(Q') \rangle_L$ in $\mathcal{C}_{\mathcal{R}}$. But $Q \approx Q'$ implies $FAM(P) \approx FAM(Q')$ in $\mathcal{C}_{\mathcal{R}}$ by Lemma 46, and therefore $\langle FAM(P) \rangle_L \trianglelefteq \langle FAM(Q) \rangle_L$ in $\mathcal{C}_{\mathcal{R}}$.
- (\Leftarrow) By Definition 44, $\exists \alpha : \alpha \approx FAM(Q) \wedge FAM(P) \subseteq \alpha$; thus by Lemma 47 $\exists [\beta] : FAM(P) \subseteq \beta \subseteq \alpha \wedge \beta \approx \alpha$ (assuming that say e_0, e_1, \dots, e_n is an enumeration of α whose initial part is an enumeration of $FAM(P)$, we can take e_{i_0}, \dots, e_{i_m} as in Lemma 47 for $[\beta]$). By Lemma 42, there is a Q' such that $FAM(Q') = \beta$ and P is an initial part of Q' . By Lemma 46, $Q \approx Q'$, hence $P \trianglelefteq Q$, therefore $\langle P \rangle_L \trianglelefteq \langle Q \rangle_L$.

Theorem 49 Let \mathcal{R} be an ASDRS and let $\mathcal{C} = \mathcal{C}_{\mathcal{R}}$ be its corresponding DEES. Then $f : \langle P \rangle_L \mapsto \langle FAM(P) \rangle_L$ is the isomorphism between $\mathcal{L}_{fin}^{\triangleleft}(\mathcal{R})$ and $\mathcal{L}_{fin}^{\triangleleft}(\mathcal{C})$, with the inverse $f^{-1} : \langle \alpha \rangle_L \mapsto \langle [ST(\alpha)] \rangle_L$. Moreover, reductions in $\langle P \rangle_L$ and $\langle FAM(P) \rangle_L$ are in one-to-one correspondence.

Proof By Lemmas 46 and 39 and Proposition 25, f and f^{-1} are well defined. The rest is immediate from Lemma 42 and Proposition 48.

A similar statement is valid when \mathcal{C} is an arbitrary DEES and $\mathcal{R} = \mathcal{R}_{\mathcal{C}}$: We only need to replace $FAM(P)$ in the definition of f with $EVN(P)$, where $EVN(P)$ is the set of events fired in P .

6 Preservation of Neededness and Stability by the Translations

This section develops a normalization-by-neededness theory for DEESs, and relates it with that in affine SDRSs, establishing a relationship between the concepts of stability intrinsic to the above models, and between neededness and stability in each model.

In order to extend the theory of Relative Normalization [9–11] from SDRSs

to DEESs, we can restrict ourselves to ASDRSs of the form $\mathcal{R}_{\mathcal{C}}$, where \mathcal{C} is a DEES: by Theorem 49, such ASDRSs contain an isomorphic copy of any ASDRS.

Definition 50 ([9]) Let \mathcal{S} be a set of terms in a DRS \mathcal{R} .

- (1) We call a redex $u \subseteq t$ \mathcal{S} -needed if $t \notin \mathcal{S}$ and at least one residual of u is contracted in any reduction from t to a term in \mathcal{S} ; we and call u \mathcal{S} -unnneeded otherwise.
- (2) We call \mathcal{S} stable⁵ if:
 - (a) \mathcal{S} is closed under reduction; and
 - (b) \mathcal{S} is closed under unnneeded expansion: for any $e \xrightarrow{u} o$ such that $e \notin \mathcal{S}$ and $o \in \mathcal{S}$, u is \mathcal{S} -needed.

All interesting sets of ‘finite results’, such as of normal forms, head-normal forms, weak-head-normal forms, etc. are stable [10]. It has been shown in [11] (a corollary of the so called Relative Normalization theorem) that, for any stable set \mathcal{S} of terms in an ASDRS \mathcal{R} and any \mathcal{S} -normalizable term $t \notin \mathcal{S}$, t contains at least one \mathcal{S} -needed redex, and repeated contraction of \mathcal{S} -needed redexes is \mathcal{S} -normalizing, which justifies the concept of stable sets.

Our translation of AZDFSs into DEESs suggests the following definition of relative neededness and stability in DEESs.

Definition 51 Let \mathcal{S}' be a set of configurations in a DEES $\mathcal{C} = (E, \leq, \triangleright, \approx)$.

- (1) We call $e \in E$ \mathcal{S}' -needed, written $NE_{\mathcal{S}'}(e)$ if, for any $\alpha \in \mathcal{S}'$, $e \in \alpha$, and call it \mathcal{S}' -unnneeded otherwise.
- (2) We call \mathcal{S}' stable if:
 - (a) \mathcal{S}' is closed under inclusion.
 - (b) \mathcal{S}' is closed under unnneeded expansion: $\alpha + e \in \mathcal{S}' \wedge \alpha \notin \mathcal{S}' \Rightarrow NE_{\mathcal{S}'}(e)$; and
 - (c) \mathcal{S}' is closed under permutation equivalence.

Note that the relative neededness concept already makes sense for PESs.

The next theorem gives two equivalent definitions of stability in DEESs, more in the spirit of the definition of stability in PESs [37], and more convenient to generalize to the case of infinite configurations.

Theorem 52 Equivalent definitions of stability of sets of finite configurations in a DEES can be obtained from Definition 51 by replacing the condition (b) with either of the following two conditions (b') or (b''):

⁵ Our definition follows [9] rather than [10,11]; in the latter papers, the definition of stability of sets allows a slightly weaker condition than closure under reduction.

- (b') \mathcal{S}' is closed under intersection.
(b'') \mathcal{S}' has a unique \subseteq -minimal element $\alpha_{\mathcal{S}'}$.

Proof We show that (i) (b) \Rightarrow (b'); (ii) (b') \Rightarrow (b'') and (iii) (a), (b'') \Rightarrow (b).

- (i) Let $\alpha, \beta \in \mathcal{S}'$, and suppose on the contrary that $\alpha \cap \beta \notin \mathcal{S}'$. Further, let $e_1, \dots, e_m, \dots, e_n$ be a left-closed enumeration of α such that e_1, \dots, e_m is a left-closed enumeration of $\alpha \cap \beta$ (such an enumeration exists since $\alpha \cap \beta$ is left-closed). Then there is $m \leq j < n$ such that $\{e_1, \dots, e_j\} \notin \mathcal{S}'$ and $\{e_1, \dots, e_{j+1}\} \in \mathcal{S}'$. By (b), e_{j+1} is \mathcal{S}' -needed, which is not possible since $e_{j+1} \notin \beta \in \mathcal{S}'$.
- (ii) If \mathcal{S}' does not have a unique \subseteq -minimal element, then it has at least two different \subseteq -minimal elements α and β (i.e., $\nexists \gamma \subset \alpha : \gamma \in \mathcal{S}'$, and similarly for β). By (b'), $\alpha \cap \beta \in \mathcal{S}'$. But $\alpha \cap \beta \subset \alpha$, contradicting the minimality of α .
- (iii) Let α, e be such that $\alpha \notin \mathcal{S}' \wedge \alpha + e \in \mathcal{S}'$. By (a) and (b''), this means that $\alpha_{\mathcal{S}'} \not\subseteq \alpha \wedge \alpha_{\mathcal{S}'} \subseteq \alpha + e$. Hence $e \in \alpha_{\mathcal{S}'}$, and therefore $\forall \delta \in \mathcal{S}' : e \in \delta$ (since $\alpha_{\mathcal{S}'} \subseteq \delta$ by (b'')).

The proof of the following proposition is routine, and therefore omitted.

Proposition 53 Let \mathcal{S}' be a set of configurations, in a DEES \mathcal{C} , closed under permutation equivalence, and let \mathcal{S} be the set of corresponding terms in the corresponding ASDRS $\mathcal{R}_{\mathcal{C}}$ (which are Lévy-equivalence classes of configurations in \mathcal{C}). Then \mathcal{S}' is stable in \mathcal{C} iff \mathcal{S} is stable in $\mathcal{R}_{\mathcal{C}}$. Further, if \mathcal{S}' is stable, an event e is \mathcal{S}' -needed in \mathcal{C} iff any e -redex is \mathcal{S} -needed in $\mathcal{R}_{\mathcal{C}}$.

We conclude this section with a theorem expressing the relationship between concepts of stability and neededness in DEESs, analogous to the Relative Normalization Theorem for ASDRSs [11]:

Theorem 54 Let \mathcal{S}' be a stable set of configurations in a DEES \mathcal{C} . Then a repeated execution of \mathcal{S}' -needed events in any configuration $\alpha \notin \mathcal{S}'$, possibly interleaved by finite sequences of \mathcal{S}' -unneded steps, eventually yields a configuration in \mathcal{S}' .

Proof Immediate from closure of \mathcal{S}' under inclusion (since \mathcal{S}' consists of finite configurations, hence there are only a finite number of \mathcal{S}' -needed events).

7 Relativized Information Domains

In this section, we introduce orderings on configurations relativized w.r.t. stable sets of results, in a DEES \mathcal{C} , and show that they form finitary prime algebraic complete lattices. Unlike the previous sections, we will now allow

for infinite configurations – (countably) infinite left-closed sets of events in \mathcal{C} . The set of finite and infinite configurations in \mathcal{C} will be denoted by $\mathcal{L}(\mathcal{C})$.

We start by some preliminary lattice theoretic definitions; we follow [37].

Definition 55 • A *complete lattice* is a partial order $\mathcal{L} = (L, \sqsubseteq)$ which has least upper bounds (joins or suprema) $\sqcup X$ and greatest lower bounds (meets or infima) $\sqcap X$ of arbitrary subsets $X \subseteq L$. We write $x \sqcup y$ and $x \sqcap y$ for the join and meet respectively of two elements $x, y \in L$.

- A *directed subset* of a partial order \mathcal{L} is a subset $Y \subseteq L$ with the property that for any finite set $X \subseteq Y$ there is an element $y \in Y$ such that $\forall x \in X. x \sqsubseteq y$.
- A *finite element* of a complete lattice is an element z with the property that, for all directed subsets Y , if $z \sqsubseteq \sqcup Y$, then there is some $y \in Y$ for which $z \sqsubseteq y$.
- A complete lattice \mathcal{L} is *algebraic* if for any element $d \in L$ the set $\{x \sqsubseteq d \mid x \text{ is finite}\}$ is directed and has least upper bound d .
- A *complete prime* of a complete lattice \mathcal{L} is an element $p \in L$ such that $p \sqsubseteq \cup X \Rightarrow \exists x \in X. p \sqsubseteq x$ for any subset $X \subseteq L$. \mathcal{L} is *prime algebraic* if, for all $x \in L$, $x = \sqcup \{p \sqsubseteq x \mid p \text{ is a complete prime}\}$.
- An algebraic lattice is *finitary* iff every finite element dominates only a finite number of elements, i.e., $\{x \mid x \sqsubseteq d\}$ is finite for every finite element d .
- A complete lattice \mathcal{L} is *distributive* if $x \sqcap (y \sqcup z) = (x \sqcap y) \sqcup (x \sqcap z)$, for all elements $x, y, z \in L$.

Theorem 56 ([37]) Let \mathcal{L} be a finitary algebraic lattice. Then \mathcal{L} is prime algebraic iff it is distributive.

We now extend some of the previously introduced concepts from finite to infinite configurations. The concepts of an α -reduction and a \mathcal{S}' -needed event generalize immediately.

Definition 57 Let $\mathcal{C} = (E, \trianglelefteq, \triangleright, \approx)$ be a DEES.

- (1) Let $\alpha \in \mathcal{L}(\mathcal{C})$. Then $\alpha \triangleright e$ iff $\exists \beta \subseteq_{fin} \alpha : \beta \triangleright e$.
- (2) Let $\alpha, \beta \in \mathcal{L}(\mathcal{C})$. Then $\alpha \approx \beta$ iff $\alpha \triangleright \beta \setminus \alpha \wedge \beta \triangleright \alpha \setminus \beta$.
- (3) We call $\mathcal{S}' \subseteq \mathcal{L}(\mathcal{C})$ *stable* if it is closed under \sqsubseteq , has a unique \sqsubseteq -minimal element $\alpha_{\mathcal{S}'}$, and is closed under permutation equivalence.

It is easy to check that Definitions 57.(1)-(2) do not conflict with the definitions of the corresponding concepts for finite configurations. Further, it is an easy consequence of Definition 57.(1) and [E1] that:

Lemma 58 Let $\alpha \triangleright e$ and $\alpha \subseteq \beta \in \mathcal{L}(\mathcal{C})$. Then $\beta \triangleright e$.

We call a reduction in a DEES \mathcal{S}' -needed-fair, or simply \mathcal{S}' -fair, if it fires at

least all \mathcal{S}' -needed events. \mathcal{S}' -fair configurations are defined similarly.

Theorem 59 Let \mathcal{S}' be a stable set of configurations in a DEES \mathcal{C} . Then its \subseteq -minimal element $\alpha_{\mathcal{S}'}$ consists of all \mathcal{S}' -needed events of \mathcal{C} . Consequently, \mathcal{S}' consists of all \mathcal{S}' -fair configurations, and any \mathcal{S}' -fair reduction generates a configuration in \mathcal{C} .

Proof Since $\alpha_{\mathcal{S}'} \in \mathcal{S}'$, all \mathcal{S}' -needed events of \mathcal{C} are in $\alpha_{\mathcal{S}'}$. If on the contrary $\alpha_{\mathcal{S}'}$ contains an \mathcal{S}' -unnneeded event e , then $\exists \beta \in \mathcal{S}' : e \notin \beta$, hence $\beta \not\subseteq \alpha_{\mathcal{S}'}$, contradicting the minimality of $\alpha_{\mathcal{S}'}$. The rest is immediate.

For example, firing all events at any state, or all \mathcal{S}' -needed events, in parallel yields an \mathcal{S}' -fair reduction since any finite configuration not yet in \mathcal{S}' can fire at least one \mathcal{S}' -needed event.

Note that the set of all *fair configurations* – configurations $\alpha \in \mathcal{C}$ such that $\forall e \in \mathcal{C} : e \notin \alpha \Rightarrow \alpha \triangleright e$ – need not be stable for every DEES \mathcal{C} . The reason is that, unlike the finite case, a configuration α need not be equivalent to its standard variant $ST_{\triangleright}(\alpha) = \{e \in \alpha \mid \alpha \not\triangleright e\}$. Indeed, consider the DEES corresponding to the ASDRS from [17] demonstrating the failure of the infinitary standardization theorem: there are infinitely many events $e_0, e'_1, e_1, e'_2, e_2, \dots$, the causal dependency is the transitive reflexive closure of dependencies $e_i < e_{i+1}, e'_{i+1}$ ($i = 0, 1, \dots$), and the erasure relation is defined by $\alpha \triangleright e'_j$ iff $e'_k \in \alpha$ for some $k > j$, $k = 2, 3, \dots$. Then any fair configuration consists of all non-erasable events e_i , and at least one infinite sequence of erasable events e'_i , and it is immediate that the set of all fair configurations does not have the \subseteq -minimal element. Note that for any fair configuration α in that DEES, $ST(\alpha)$ consists of all non-erasable events e_i , and $\alpha \not\approx ST(\alpha)$ since $ST(\alpha) \not\triangleright e'_k$ for any k . Thus $ST(\alpha) = ST(\beta)$ does not imply $\alpha \approx \beta$ as $ST(ST(\alpha)) = ST(\alpha)$. However, the converse is always true:

Lemma 60 Let $\alpha, \beta \in \mathcal{L}(\mathcal{C})$. Then $\alpha \approx \beta$ implies $ST(\alpha) = ST(\beta)$.

Proof By Definition 57.(2), $\alpha \triangleright \beta \setminus \alpha \cap \beta$ and $\beta \triangleright \alpha \setminus \alpha \cap \beta$. Hence $ST(\alpha), ST(\beta) \subseteq \alpha \cap \beta$. Suppose on the contrary that $\exists e \in ST(\alpha) \setminus ST(\beta)$. Then $\alpha \not\triangleright e$ and $\beta \triangleright e$, i.e., $\exists \beta_0 \subseteq_{fin} \beta : \beta_0 \triangleright e$, implying by Lemma 58 that $\alpha \cap \beta \not\triangleright e$ and $\beta' = \beta_0 \cap (\alpha \cap \beta) \not\triangleright e$. Therefore, by Proposition 21, $\exists e' \in \beta_0 \setminus \beta' : \beta' \not\triangleright e'$. Thus $\alpha \not\triangleright e'$ by Lemma 58 (since $\beta' \subseteq \alpha \cap \beta \subseteq \alpha$) and $e' \in \beta \setminus \alpha \cap \beta$, contradicting $\alpha \triangleright \beta \setminus \alpha \cap \beta$.

Definition 61 Let $\alpha, \beta \in \mathcal{L}(\mathcal{C})$, and let \mathcal{S}' be a stable set of configurations in \mathcal{C} . We define $\alpha \trianglelefteq_{\mathcal{S}'} \beta$ if $[\alpha]_{\mathcal{S}'} \subseteq [\beta]_{\mathcal{S}'}$, where $[\alpha]_{\mathcal{S}'}$ denotes the set of \mathcal{S}' -needed events of α . We write $\alpha \approx_{\mathcal{S}'} \beta$ iff $\alpha \trianglelefteq_{\mathcal{S}'} \beta$ and $\beta \trianglelefteq_{\mathcal{S}'} \alpha$, i.e., $[\alpha]_{\mathcal{S}'} = [\beta]_{\mathcal{S}'}$. We call $\trianglelefteq_{\mathcal{S}'}$ a *stable embedding* or *\mathcal{S}' -embedding*, and call $\approx_{\mathcal{S}'}$ a *stable equivalence* or *\mathcal{S}' -equivalence*. $\langle \alpha \rangle_{\mathcal{S}'}$ will denote the $\approx_{\mathcal{S}'}$ -equivalence class of α . We write $\langle \alpha \rangle_{\mathcal{S}'} \trianglelefteq_{\mathcal{S}'} \langle \beta \rangle_{\mathcal{S}'}$ if $\alpha \trianglelefteq_{\mathcal{S}'} \beta$. Finally, we define $\mathcal{L}^{\approx_{\mathcal{S}'}}(\mathcal{C}) = \mathcal{L}(\mathcal{C}) / \approx_{\mathcal{S}'} = \{\langle \alpha \rangle_{\mathcal{S}'} \mid \alpha \in$

$\mathcal{L}(\mathcal{C})\}$ and $\mathcal{L}^{\triangleleft_{\mathcal{S}'}}(\mathcal{C}) = (\mathcal{L}^{\approx_{\mathcal{S}'}}(\mathcal{C}), \triangleleft_{\mathcal{S}'})$.

Note that $\approx_{\mathcal{S}'}$ does not satisfy [P0], although it does satisfy the other permutation axioms [P1] - [P3]. This means that for permutation equivalence ‘*progress in any direction*’ is equally important, while for a stable equivalence $\approx_{\mathcal{S}'}$ only ‘*progress towards \mathcal{S}'* ’ matters. Actually, \approx is a special case of $\approx_{\mathcal{S}'}$: the two coincide when \mathcal{C} is a PES and \mathcal{S}' is the set of all events.

Definition 62 Let \mathcal{S}' be a stable set of configurations in a DEES \mathcal{C} . The meet and join operations $\sqcup_{\mathcal{S}'}, \sqcap_{\mathcal{S}'}$ on $\mathcal{L}^{\triangleleft_{\mathcal{S}'}}(\mathcal{C})$ are defined by:

- $\alpha \sqcup_{\mathcal{S}'} \beta = [\alpha]_{\mathcal{S}'} \cup [\beta]_{\mathcal{S}'}$ and $\langle \alpha \rangle_{\mathcal{S}'} \sqcup_{\mathcal{S}'} \langle \beta \rangle_{\mathcal{S}'} = \langle \alpha \sqcup_{\mathcal{S}'} \beta \rangle_{\mathcal{S}'}$;
- $\alpha \sqcap_{\mathcal{S}'} \beta = [\alpha]_{\mathcal{S}'} \cap [\beta]_{\mathcal{S}'}$ and $\langle \alpha \rangle_{\mathcal{S}'} \sqcap_{\mathcal{S}'} \langle \beta \rangle_{\mathcal{S}'} = \langle \alpha \sqcap_{\mathcal{S}'} \beta \rangle_{\mathcal{S}'}$;

It is immediate from the definition of $\approx_{\mathcal{S}'}$ that the above definition is correct.

Lemma 63 Let $\beta \triangleright e$, in a DEES \mathcal{C} . Then $\beta \approx \beta \setminus [e]^{\geq}$.

Proof If β is finite, then the lemma follows from Proposition 21. Otherwise, we have from Definition 57.(1)-(2) and [E3] that $\beta \triangleright e \Rightarrow \exists \beta_0 \subseteq_{fin} \beta : \beta_0 \triangleright e \Rightarrow \beta_0 \triangleright [e]^{\geq} \Rightarrow \beta \triangleright [e]^{\geq} \Rightarrow \beta \approx \beta \setminus [e]^{\geq}$.

Lemma 64 Let \mathcal{S}' be a stable set of configurations in a DEES \mathcal{C} , and let e be \mathcal{S}' -needed. Then e is non-erasable: $\forall \alpha \in \mathcal{L}(\mathcal{C}) : \alpha \not\triangleright e$.

Proof Suppose on the contrary that $\exists \alpha \in \mathcal{L}(\mathcal{C}) : \alpha \triangleright e$. We can assume that $\alpha \in \mathcal{L}_{fin}(\mathcal{C})$. By [E3] and [E2], $\alpha \setminus [e]^{\geq} \triangleright e$. But since e is \mathcal{S}' -needed, $\alpha \setminus [e]^{\geq} \notin \mathcal{S}'$. Hence $\beta = \alpha_{\mathcal{S}'} \cup (\alpha \setminus [e]^{\geq}) \in \mathcal{S}'$ by closure of \mathcal{S}' under inclusion. But $\beta \triangleright e$ by Lemma 58, and $\beta \approx \beta \setminus [e]^{\geq}$ by Lemma 63. Thus, by closure of \mathcal{S}' under permutation equivalence, $\beta \setminus [e]^{\geq} \in \mathcal{S}'$, and $e \notin \beta \setminus [e]^{\geq}$, contradicting \mathcal{S}' -neededness of e .

The following proposition shows that $\approx \subseteq \approx_{\mathcal{S}'}$, which agrees with our intuition that a concurrent semantics must not discriminate between permutation equivalent configurations/reductions.

Proposition 65 Let α, β be configurations in a DEES \mathcal{C} , and let \mathcal{S}' be a stable set of configurations in \mathcal{C} . Then $\alpha \approx \beta$ implies $\alpha \approx_{\mathcal{S}'} \beta$.

Proof By Definition 57.(1), $\alpha \triangleright \beta \setminus \alpha$ and $\beta \triangleright \alpha \setminus \beta$. Hence, by Lemma 64, $[\alpha]_{\mathcal{S}'}, [\beta]_{\mathcal{S}'} \subseteq \alpha \cap \beta$, and by Theorem 59, $[\alpha]_{\mathcal{S}'} = \alpha \cap \alpha_{\mathcal{S}'} = (\alpha \cap \beta) \cap \alpha_{\mathcal{S}'} = \beta \cap \alpha_{\mathcal{S}'} = [\beta]_{\mathcal{S}'}$.

Theorem 66 Let \mathcal{S}' be a stable set of configurations in a DEES \mathcal{C} . Then $\mathcal{L}^{\triangleleft_{\mathcal{S}'}}(\mathcal{C}) = (\mathcal{L}^{\approx_{\mathcal{S}'}}(\mathcal{C}), \triangleleft_{\mathcal{S}'})$ is a finitary prime algebraic complete lattice.

Proof Immediate from Definition 62 and Theorem 59, since the set of subsets

of any non-empty set (in our case, $\alpha_{S'}$) ordered by the subset relation is a finitary prime algebraic complete lattice with intersection and union as meet and join operations [37].

In an ASDRS \mathcal{R} , stability of a set \mathcal{S} of initial (finite or infinite) reductions can be defined via stability of the corresponding set \mathcal{S}' of configurations in $\mathcal{C}_{\mathcal{R}}$. For any initial reductions P, Q , and any stable set of reductions \mathcal{S} , in an ASDRS \mathcal{R} , we can define $P \trianglelefteq_{\mathcal{S}} Q$ iff $FAM_{\mathcal{S}}(P) \trianglelefteq_{\mathcal{S}'} FAM_{\mathcal{S}}(Q)$ in $\mathcal{C}_{\mathcal{R}}$, where $FAM_{\mathcal{S}}(P)$ denotes the set of \mathcal{S} -needed families contracted in P (and hence coincides with the set of \mathcal{S}' -needed events in the configuration corresponding to P). Further, we define $P \approx_{\mathcal{S}} Q$ iff $P \trianglelefteq_{\mathcal{S}} Q$ and $Q \trianglelefteq_{\mathcal{S}} P$. We call $\trianglelefteq_{\mathcal{S}}$ and $\approx_{\mathcal{S}}$ a *stable embedding* or *\mathcal{S} -embedding* and a *stable equivalence* or *\mathcal{S} -equivalence*, respectively. $\langle P \rangle_{\mathcal{S}}$ denotes the $\approx_{\mathcal{S}}$ -equivalence class of P . We write $\langle P \rangle_{\mathcal{S}} \trianglelefteq_{\mathcal{S}} \langle Q \rangle_{\mathcal{S}}$ if $P \trianglelefteq_{\mathcal{S}} Q$. Now, if $\mathcal{L}(\mathcal{R})$ is the set of (finite or infinite) initial reductions in \mathcal{R} , we can define $\mathcal{L}^{\approx_{\mathcal{S}}}(\mathcal{R}) = \mathcal{L}(\mathcal{R}) / \approx_{\mathcal{S}} = \{\langle P \rangle_{\mathcal{S}} \mid P \in \mathcal{L}(\mathcal{R})\}$ and $\mathcal{L}^{\trianglelefteq_{\mathcal{S}}}(\mathcal{R}) = (\mathcal{L}^{\approx_{\mathcal{S}}}(\mathcal{R}), \trianglelefteq_{\mathcal{S}})$.

Lemma 67 Let $\alpha \in \mathcal{L}(\mathcal{C})$. Then $[\alpha]_{\mathcal{S}'} \in \mathcal{L}(\mathcal{C})$. Moreover, $[\alpha]_{\mathcal{S}'}$ is standard, and any of its left-closed enumerations is a reduction.

Proof Let $e, e' \in \alpha$, let $e < e'$ and let e' be \mathcal{S}' -needed. We need to show that e is \mathcal{S}' -needed too. By Theorem 59, $e' \in \alpha_{\mathcal{S}'} \in \mathcal{L}(\mathcal{C})$, implying that $e \in \alpha_{\mathcal{S}'}$. Thus $[\alpha]_{\mathcal{S}'} \in \mathcal{L}(\mathcal{C})$ ($[\alpha]_{\mathcal{S}'}$ is left-closed). The rest is immediate from Lemma 64.

Theorem 68 Let \mathcal{R} be an ASDRS and let $\mathcal{C} = \mathcal{C}_{\mathcal{R}}$ be its corresponding DEES. Further, let \mathcal{S}' be a set of configurations in \mathcal{C} , and let \mathcal{S} be the set of corresponding reductions in \mathcal{R} . Then $f : \langle P \rangle_{\mathcal{S}} \rightarrow \langle FAM(P) \rangle_{\mathcal{S}'}$ is an isomorphism between $\mathcal{L}^{\trianglelefteq_{\mathcal{S}}}(\mathcal{R})$ and $\mathcal{L}^{\trianglelefteq_{\mathcal{S}'}}(\mathcal{C})$, with the inverse $f^{-1} : \langle \alpha \rangle_{\mathcal{S}'} \rightarrow \langle [[\alpha]_{\mathcal{S}'}] \rangle_{\mathcal{S}}$.

Proof By Lemma 67, $[[\alpha]_{\mathcal{S}'}]$ and hence f^{-1} are well defined. The rest is immediate from the definitions above.

Hence we can define on $\mathcal{L}^{\trianglelefteq_{\mathcal{S}}}(\mathcal{R})$ that $\langle P \rangle_{\mathcal{S}} \sqcup_{\mathcal{S}} \langle Q \rangle_{\mathcal{S}} = f^{-1}(f(\langle P \rangle_{\mathcal{S}}) \sqcup_{\mathcal{S}'} f(\langle Q \rangle_{\mathcal{S}}))$ and $\langle P \rangle_{\mathcal{S}} \sqcap_{\mathcal{S}} \langle Q \rangle_{\mathcal{S}} = f^{-1}(f(\langle P \rangle_{\mathcal{S}}) \sqcap_{\mathcal{S}'} f(\langle Q \rangle_{\mathcal{S}}))$, and we get from Theorem 66 that $\mathcal{L}^{\trianglelefteq_{\mathcal{S}}}(\mathcal{R})$ with $\sqcup_{\mathcal{S}}$ and $\sqcap_{\mathcal{S}}$ as join and meet is a finitary prime algebraic domain. It is not difficult to define stability of a set of reductions \mathcal{S} , as well as the relations $\trianglelefteq_{\mathcal{S}}, \sqcup_{\mathcal{S}}$ and $\sqcap_{\mathcal{S}}$, on (corresponding equivalence classes of) reductions in \mathcal{R} directly.

8 Conclusions

We have established an equivalence between deterministic stable operational (ASDRS) and domain-theoretic (DEES) models of computation, and based on this, constructed a fully adequate event style concurrent semantics for orthog-

onal rewrite systems. The correspondence between Event Structures and other models of concurrency, as well as with denotational semantics, are well studied [37,38], and we hope that our results contribute to better understanding of the relationship between operational and denotational semantics of sequential and concurrent languages.

We think that our axiomatization of permutation equivalence is interesting on its own right from the poset/lattice theoretic point of view. Definitions of permutation equivalence without using a residual calculus have been studied before [25], but they still use the syntax of the rewrite system essentially (axioms depend on the syntax).

Although we have restricted attention to orthogonal rewriting, it would be interesting to extend our axiomatization to partial orders with conflicts, to enable modelling of non-deterministic calculi as well.

Acknowledgments We thank R. Kennaway for useful comments.

References

- [1] A. Asperti and S. Guerrini, *The Optimal Implementation of Functional Programming Languages*, Cambridge Tracts in Theoretical Computer Science, vol. 45, (Cambridge University Press, 1998).
- [2] A. Asperti and H.G. Mairson, Parallel beta reduction is not elementary recursive, in: *Proc. 15th ACM Symposium on Principles of Programming Languages* (1998) 303-315.
- [3] H. P. Barendregt, *The Lambda Calculus, its Syntax and Semantics* (North-Holland, Amsterdam, 1984).
- [4] G. Berry and J.-J. Lévy, Minimal and optimal computations of recursive programs, *J. of the Association for Computing Machinery* **26** (1979) 148-175.
- [5] G. Boudol, Computational semantics of term rewriting systems. in: M. Nivat, J. C. Reynolds, eds., *Algebraic methods in semantics* (Cambridge University Press, 1985) 169-236.
- [6] G. Boudol and I. Castellani, A non-interleaving semantics for CCS based on proved transitions, *Fundamenta Informaticae* **11(4)** (1988) 433-452.
- [7] D. Clark and J.R. Kennaway, Event structures and non-orthogonal term graph rewriting, *Mathematical Structures in Computer Science* **6** (1996) 545-578.
- [8] A. Corradini, F. Gadduchi and U. Montanari, Relating two categorical models of term rewriting, in: *Proc. RTA '95*, Lecture Notes in Computer Science, vol. 914 (Springer, Berlin, 1995) 225-240.

- [9] J.R.W. Glauert, J.R. Kennaway and Z. Khasidashvili, Stable results and relative normalization, *Journal of Logic and Computation*, **10(3)** (2000). Special Issue on Type Theory and Term Rewriting, F. Kamareddine and J.W. Klop, eds. Oxford University Press.
- [10] J.R.W. Glauert and Z. Khasidashvili, Relative normalization in orthogonal expression reduction systems, in: *Proc. CTRS'94*, Lecture Notes in Computer Science, vol. 968 (Springer, Berlin, 1994) 144-165.
- [11] J.R.W. Glauert and Z. Khasidashvili, Relative normalization in deterministic residual structures. in: *Proc. CAAP'96*, Lecture Notes in Computer Science, vol. 1059 (Springer, Berlin, 1996) 180-195.
- [12] G. Gonthier, J.-J. Lévy and P.-A. Melliès, An abstract standardisation theorem, in: *Proc. 7th IEEE Symp. on Logic in Computer Science* (1992) 72-81.
- [13] R.J. Hindley, An abstract form of the Church-Rosser theorem I, *Journal of Symbolic Logic* **34(4)** (1969) 545-560.
- [14] G. Huet and J.-J. Lévy, Computations in Orthogonal Rewriting Systems, in: J.-L. Lassez and G. Plotkin, eds., *Computational Logic, Essays in Honor of Alan Robinson* (MIT Press, 1991) 394-443.
- [15] V. Kathail, Optimal interpreters for lambda-calculus based functional languages, Ph.D. Thesis, MIT, 1990.
- [16] J.R. Kennaway, J.W. Klop, M.R. Sleep and F.-J. de Vries, Event structures and orthogonal term graph rewriting, in: M.R. Sleep, M.J. Plasmeijer, M.C.J.D. van Eekelen, eds., *Term Graph Rewriting: Theory and Practice* (John Wiley, 1993) 141-156.
- [17] Z. Khasidashvili and J.R.W. Glauert, Discrete normalization and standardization in deterministic residual structures, in: *Proc. ALP'96*, Lecture Notes in Computer Science, vol. 1139 (Springer, Berlin, 1996) 135-149.
- [18] Z. Khasidashvili and J.R.W. Glauert, Zig-zag, extraction and separable families in non-duplicating stable deterministic residual structures, Technical Report IR-420, Free University, 1997.
- [19] Z. Khasidashvili and J.R.W. Glauert, Relating conflict-free stable transition and event models (extended abstract), in: *Proc. MFCS'97*, Lecture Notes in Computer Science, vol. 1295, (Springer, Berlin, 1997) 269-278.
- [20] Z. Khasidashvili and J.R.W. Glauert, An abstract concept of optimal implementation, Technical Report SYS-C97-01, UEA Norwich, 1997.
- [21] J. W. Klop, *Combinatory reduction systems* (Mathematical Centre Tracts n. 127, Amsterdam, 1980).
- [22] J. W. Klop, Term rewriting systems in: S. Abramsky, D. Gabbay and T. Maibaum, eds., *Handbook of Logic in Computer Science, vol. 2I* (Oxford University Press, 1992) 1-116.

- [23] J. Lamping, An algorithm for optimal lambda calculus reduction, in: *Proc. 7th ACM Symp. on Principles of Programming Languages* (1990) 6-30.
- [24] C. Laneve, Distributive evaluations of λ -calculus, *Fundamenta Informaticae* **20(4)** (1994) 333-352.
- [25] C. Laneve and U. Montanari, Axiomatizing permutation equivalence, *Mathematical Structures in Computer Science* **6** (1994) 219 - 249.
- [26] J.-J. Lévy, Réductions correctes et optimales dans le lambda-calcul, Thèse de l'Université de Paris VII, 1978.
- [27] J.-J. Lévy, Optimal reductions in the Lambda-calculus, in: J. P. Seldin, J. R. Hindley, eds., *To H. B. Curry: Essays on Combinatory Logic, Lambda-calculus and Formalism* (Academic Press, 1980) 159-192.
- [28] L. Maranget, Optimal derivations in weak λ -calculi and in orthogonal term rewriting systems, in: *Proc. 18th ACM Symp. on Principles of Programming Languages* (1991) 255-269.
- [29] P.-A. Melliès, Description Abstraite des Systèmes de Réécriture, Thèse de l'Université Paris 7, 1996.
- [30] P.-A. Melliès, A factorization theorem in rewriting theory, in; *Proc. CTCS'97*, Lecture Notes in Computer Science, vol. 1290 (Springer, Berlin, 1997) 49-68.
- [31] R. Milner, *Communication and Concurrency* (Prentice Hall, 1989).
- [32] M. Nielsen, G. Plotkin and G. Winskel, Petri nets, event structures and domains, Part 1, *Theoret. Comput. Sci.* **13** (1981) 85-108.
- [33] D. Sangiorgi, Expressing mobility in process algebras: first-order and higher-order paradigms, Ph.D. Thesis, Edinburgh University, 1993.
- [34] G. Schied, On relating rewrite systems and graph grammars to event structures, in: *Proc. GTCS'94*, Lecture Notes in Computer Science, vol. 776, (Springer, Berlin, 1994) 326-340.
- [35] E.W. Stark, Concurrent transition systems, *Theoret. Comput. Sci.* **64(3)** (1989) 221-269.
- [36] Winskel G. Events in Computation, Ph.D. Thesis, Edinburgh University, 1980.
- [37] Winskel G. An introduction to event structures, In Proc. 'Linear time, branching time and partial order in logics and models of concurrency', Lecture Notes in Computer Science, vol. 354, (Springer, Berlin, 1989) 364-397. Extended version appears as Cambridge University Computer Laboratory Report n. 95, 1986.
- [38] G. Winskel and M. Nielsen, Models for concurrency, in: S. Abramsky, D. Gabbay, and T. Maibaum, eds., *Handbook of Logic in Computer Science*, vol. 4 (Oxford University Press, 1995) 1-148.